



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5: C07K 13/00, C12P 21/00, C12N 5/00, 15/00	A1	(11) International Publication Number: WO 94/21679 (43) International Publication Date: 29 September 1994 (29.09.94)
(21) International Application Number: PCT/US94/01957 (22) International Filing Date: 25 February 1994 (25.02.94) (30) Priority Data: 038,769 25 March 1993 (25.03.93) US (60) Parent Application or Grant (63) Related by Continuation US 038,769 (CIP) Filed on 25 March 1993 (25.03.93) (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): KENDALL, Richard, L. [US/US]; 5201 Stonehedge Road, Edison, NJ 08820 (US). THOMAS, Kenneth, A., Jr. [US/US]; 245 Washington Avenue, Chatham Borough, NJ 07928 (US). (74) Agent: WALLEN, John, W., III; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: INHIBITOR OF VASCULAR ENDOTHELIAL CELL GROWTH FACTOR		
(57) Abstract <p>The vascular endothelial cell growth factor (VEGF) inhibitors of the present invention are naturally occurring or recombinantly engineered soluble forms with or without a C-terminal transmembrane region of the receptor for VEGF, a very selective growth factor for endothelial cells. The soluble forms of the receptors will bind the growth factor with high affinity but do not result in signal transduction. These soluble forms of the receptor bind VEGF and inhibit its function.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5

- 1 -

10 TITLE OF THE DISCLOSURE
INHIBITOR OF VASCULAR ENDOTHELIAL CELL GROWTH FACTOR

BACKGROUND OF THE DISCLOSURE

15 Recently a new class of cell-derived dimeric
mitogens with selectivity for vascular endothelial
cells has been identified and designated vascular
endothelial cell growth factor (VEGF). VEGF has been
purified from conditioned growth media of rat glioma
cells [Conn et al., (1990), Proc. Natl. Acad. Sci.
20 U.S.A., 87, pp 2628-2632]; and conditioned growth media
of bovine pituitary folliculo stellate cells [Ferrara
and Henzel, (1989), Biochem. Biophys. Res. Comm., 161,
pp. 851-858; Gozpadorowicz et al., (1989), Proc. Natl.
Acad. Sci. U.S.A., 86, pp. 7311-7315] and conditioned
25 growth medium from human U937 cells [Connolly, D. T. et
al. (1989), Science, 246, pp. 1309-1312]. VEGF is a
dimer with an apparent molecular mass of about 46 kDa
with each subunit having an apparent molecular mass of
about 23 kDa.

30

- 2 -

VEGF has some structural similarities to platelet derived growth factor (PDGF), which is a mitogen for connective tissue cells but not mitogenic for vascular endothelial cells from large vessels.

The membrane-bound tyrosine kinase receptor, known as FLT, was shown to be a VEGF receptor [DeVries, C. *et al.*, (1992), *Science*, 255, pp.989-991]. The FLT receptor specifically binds VEGF which induces mitogenesis. Another form of the VEGF receptor, designated KDR, is also known to bind VEGF and induce mitogenesis. The partial cDNA sequence and nearly full length protein sequence of KDR is known as well [Terman, B.I. *et al.*, (1991) *Oncogene* 6, pp. 1677-1683; Terman, B.I. *et al.*, (1992) *Biochem. Biophys. Res. Comm.* 187, pp. 1579-1586].

Persistent angiogenesis may cause or exacerbate certain diseases such as psoriasis, rheumatoid arthritis, hemangiomas, angiofibromas, diabetic retinopathy and neovascular glaucoma. An inhibitor of VEGF activity would be useful as a treatment for such diseases and other VEGF-induced pathological angiogenesis and vascular permeability conditions, such as tumor vascularization.

SUMMARY OF THE DISCLOSURE

A naturally-occurring FLT messenger RNA (mRNA) was identified and cloned from vascular endothelial cells. This mRNA is shown to encode most of the extracellular, or soluble, portion of the VEGF receptor, FLT. Soluble receptor molecules including

- 3 -

forms containing a C-terminal transmembrane region are also recombinantly engineered for this and other VEGF receptors. These soluble receptors, comprising
5 truncated and modified forms are expressed in recombinant host cells and have VEGF binding properties. The soluble receptor proteins are useful as inhibitors of VEGF activity since they will bind
10 available VEGF preventing it from activating its functional receptors on vascular endothelial cells and could form non-functional heterodimers with full-length membrane anchored VEGF receptors.

BRIEF DESCRIPTION OF THE DRAWINGS

15

Figure 1 - A schematic diagram of full length VEGF receptors (FLT and KDR), the soluble VEGF receptors (sVEGF-RI and
20 sVEGF-RII) and the soluble receptors containing the C-terminal transmembrane region (sVEGF-RTMI and sVEGF-RTMII) are shown with the protein domains of each.

20

25

Figure 2 - The DNA sequence of the sVEGF-RI soluble VEGF receptor/VEGF inhibitor is shown.

30

Figure 3 - The amino acid sequence of the sVEGF-RI soluble VEGF receptor/VEGF inhibitor is shown.

Figure 4 - Demonstration that recombinant host cells express sVEGF-RI is shown by

- 4 -

the formation of high molecular weight complexes of sVEGF-RI and [^{125}I]VEGF and separated by size exclusion chromatography.

5

Figure 5 - A 12.5% polyacrylamide electrophoretic gel is shown which demonstrates the high degree of purity obtained for sVEGF-RI.

10

Figure 6 - Cross-linked products of sVEGF-RI and [^{125}I]VEGF are shown at about 145 kDa, and at about 245 kDa.

15

Figure 7A and 7B - Analysis of VEGF binding to sVEGF-RI (A) and corresponding Scatchard plot (B).

20

Figure 8 - Inhibition of [^{125}I]VEGF binding to HUVECs by sVEGF-RI is demonstrated.

Figure 9 - Inhibition of VEGF-mediated mitogenesis on HUVECs is shown using sVEGF-RI.

25

Figure 10 - The nucleotide sequence encoding sVEGF-RII is shown.

30

Figure 11 - The amino acid sequence for sVEGF-RII is shown.

- 5 -

Figure 12 - The nucleotide sequence encoding sVEGF-RTMII is shown.

5 Figure 13 - The amino acid sequence for sVEGF-RTMII is shown.

Figure 14 - The nucleotide sequence encoding sVEGF-RTMI is shown.

10 Figure 15 - The amino acid sequence for sVEGF-RTMI is shown.

Figure 16 - A diagram of pmFLT is shown.

15 Figure 17 - A diagram of pKDRA is shown.

DETAILED DESCRIPTION OF THE DISCLOSURE

The present invention relates to cDNA
20 encoding a soluble VEGF receptor protein (sVEGF-R)
which is isolated from VEGF receptor producing cells or
is recombinantly engineered from VEGF receptor-encoding
DNA. sVEGF-R, as used herein, refers to a protein
which can specifically bind to a vascular endothelial
25 cell growth factor without stimulating mitogenesis of
vascular endothelial cells.

The amino acid sequence of FLT is known,
[Shibuya, M. *et al.*, (1990), *Oncogene*, 5, pp.519-524]
and corresponds to the full length cell-associated VEGF
30 tyrosine kinase receptor. Other VEGF receptors are
known to exist. Other known VEGF receptors include,

- 6 -

but are not limited to KDR [Terman (1991), supra., and Terman (1992), supra.]. Mammalian cells capable of producing FLT, KDR and other VEGF receptors include, but are not limited to, vascular endothelial cells. Mammalian cell lines which produce FLT or KDR and other VEGF receptors include, but are not limited to, human endothelial cells. The preferred cells for the present invention include human umbilical vein endothelial cells (HUVEC).

Other cells and cell lines may also be suitable for use to isolate sVEGF-R cDNA. Selection of suitable cells may be done by screening for sVEGF-R binding activity on cell surfaces, in cell extracts or conditioned medium or by screening for gene expression by PCR or hybridization. Methods for detecting soluble receptor activity are well known in the art [Duan, D-S. R. *et al.*, (1991) J.Biol.Chem., 266, pp.413-418] and measure the binding of labelled VEGF. Cells which possess VEGF binding activity in this assay may be suitable for the isolation of sVEGF-R cDNA.

Full length FLT producing cells such as human HUVEC cells (American Type Culture Collection, ATCC CRL 1730) [Hoshi, H. and McKeehan, W.L., Proc. Natl. Acad. Sci. U.S.A., (1984) 81, pp. 6413-6417] are grown according to the recommended culture conditions of the ATCC. Full length FLT, and KDR VEGF receptors as well as extracellular region (sVEGF-RI and sVEGF-RII) and extracellular region plus transmembrane region forms (sVEGF-RTMI and sVEGF-RTMII) are shown in Figure 1. The full length receptor has an extracellular ligand

- 7 -

binding region composed of about seven immunoglobulin-like domains, a membrane spanning sequence (transmembrane domain) and intracellular tyrosine kinase domains. The inhibitory forms of this receptor, which are the subject of the present invention, are also shown in Figure 1 and lack the intracellular kinase domains, and for some inhibitors, the transmembrane sequence and the C-terminal most Ig-like extracellular domain.

Any of a variety of procedures may be used to molecularly clone sVEGF-R cDNA. These methods include, but are not limited to, direct functional expression of the sVEGF-R gene following the construction of an sVEGF-R-containing cDNA library in an appropriate expression vector system.

Another method is to screen a sVEGF-R-containing cDNA library constructed in a bacteriophage or plasmid shuttle vector with a labelled oligonucleotide probe designed from the predicted amino acid sequence of sVEGF-R. The preferred method consists of screening a sVEGF-R-containing cDNA library constructed in a bacteriophage or plasmid shuttle vector with a partial cDNA encoding at least part of the full length FLT protein. This partial cDNA is obtained by the specific PCR amplification of sVEGF-R DNA fragments through the design of oligonucleotide primers from the known sequence of the full length FLT-encoding DNA.

It is readily apparent to those skilled in the art that other types of libraries, as well as

- 8 -

libraries constructed from other cells or cell types, may be useful for isolating sVEGF-R-encoding DNA.

Other types of libraries include, but are not limited to, cDNA libraries derived from other cells or cell lines other than HUVECs and genomic DNA libraries.

It is readily apparent to those skilled in the art that suitable cDNA libraries may be prepared from cells or cell lines which have sVEGF-R activity.

The selection of cells or cell lines for use in preparing a cDNA library to isolate sVEGF-R cDNA may be done by first measuring secreted sVEGF-R activity using the VEGF binding assay described fully herein.

Preparation of cDNA libraries can be performed by standard techniques well known in the art. Well known cDNA library construction techniques can be found for example, in Maniatis, T., Fritsch, E.F., Sambrook, J., Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1982).

It is also readily apparent to those skilled in the art that DNA encoding sVEGF-R may also be isolated from a suitable genomic DNA library.

Construction of genomic DNA libraries can be performed by standard techniques well known in the art. Well known genomic DNA library construction techniques can be found in Maniatis, T., Fritsch, E.F., Sambrook, J. in Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1982).

Another means of obtaining sVEGF-R molecules is to recombinantly engineer them from DNA encoding the

- 9 -

partial or complete amino acid sequence of a VEGF receptor. Examples of other VEGF receptors include, but are not limited to, KDR. Using recombinant DNA techniques, DNA molecules are constructed which encode at least a portion of the VEGF receptor capable of binding VEGF without stimulating mitogenesis. Standard recombinant DNA techniques are used such as those found in Maniatis, *et al.*, *supra*.

Using one of the preferred methods of the present invention, cDNA clones encoding sVEGF-R are isolated in a two-stage approach employing polymerase chain reaction (PCR) based technology and cDNA library screening. In the first stage, DNA oligonucleotides derived from the extracellular domain sequence information from the known full length FLT, KDR or other VEGF receptor is used to design degenerate oligonucleotide primers for the amplification of sVEGF-R-specific DNA fragments. In the second stage, these fragments are cloned to serve as probes for the isolation of complete sVEGF-R cDNA from a commercially available lambda gt10 cDNA library (Clontech) derived from HUVEC cells (ATCC CRL 1730).

These PCR derived products were used as hybridization probes for screening a lambda gt10 cDNA library derived from HUVECs (Clontech). Plating and plaque lifts of the library were performed by standard methods (T. Maniatis, E.F. Fritsch, J. Sambrook, Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1982). The probes were random-primed labelled with

- 10 -

32P-dCTP to high specific activity and a separate screening of the library (1×10^6 plaques per screen) was conducted with each probe. The probes were
5 added to hybridization buffer (50% formamide, 5X Denhardt's, 6X SSC (1X SSC = 0.15 M NaCl, 0.015 M $\text{Na}_3\text{citrate} \cdot 2\text{H}_2\text{O}$, pH 7.0), 0.1% SDS, 100 $\mu\text{g/ml}$ salmon sperm DNA) at 1×10^6 cpm/ml.

Four positively hybridizing phage were
10 detected using the flt-specific probe. These positively hybridizing phage were observed to be less than full length flt.

Two flt cDNA clones of about 2.0 kb and 2.7 kb in length were subcloned into pGEM vectors (Promega)
15 and bi-directionally sequenced in their entirety by the chain termination method (Sanger *et al.*, (1977) P.N.A.S. USA, 74, pp. 5463-5467,) and shown to contain a single open reading frame of about 569 amino acids. Sequence analysis demonstrated that a portion of the 5'
20 flt coding region was missing from these clones. The remainder of the 5' end was cloned using PCR and combined with the DNA of the clones lacking the 5' end to yield a single open reading frame encoding about 687 amino acids.

25 The sequence for the cDNA encoding flt-derived sVEGF-RI is shown in Table 1, and was identified in clones 7 and 11. The deduced amino acid sequence of sVEGF-RI from the cloned cDNA is shown in Table 2. Inspection of the deduced amino acid sequence
30 reveals the presence of a single, large open reading frame of 687 amino acids. By comparison with amino

- 11 -

acid sequence of the full length FLT VEGF receptor, 31 amino acids are encoded at the C-terminal end of the cDNA which are different from those of FLT.

5 Using another of the preferred methods of the present invention, DNA encoding sVEGF-R is constructed from a DNA sequence encoding a VEGF receptor. For purposes of illustration, DNA encoding the VEGF receptor known as KDR was utilized. Using the receptor
10 DNA sequence, a DNA molecule is constructed which encodes the extracellular domain of the receptor, or the VEGF binding domain only and is denoted sVEGF-RII. Restriction endonuclease cleavage sites are identified within the receptor DNA and can be utilized directly to
15 excise the extracellular-encoding portion. In addition, PCR techniques as described above may be utilized to produce the desired portion of DNA. It is readily apparent to those skilled in the art that other techniques, which are standard in the art, may be
20 utilized to produce sVEGF-R molecules in a manner analogous to those described above. Such techniques are found, for example, in Maniatis *et al.*, *supra*.

Additional truncated forms of the VEGF receptor are constructed which contain the
25 transmembrane region. Retention of the transmembrane may facilitate orientation of the inhibitor molecule at the target cell surface. Examples of transmembrane region containing inhibitor molecules include but are not limited to those shown in Figure 1. sVEGF-RTMI and
30 sVEGF-RTMII, as shown in Figure 1, are FLT-related and KDR-related, respectively, transmembrane region

- 12 -

containing receptor inhibitors. Construction of transmembrane region containing molecules, such as sVEGF-RTMI and sVEGF-RTMII, is done by standard techniques known in the art including but not limited to utilizing convenient restriction endonuclease cleavage sites or PCR techniques as described herein. It is readily understood by those skilled in the art that various forms of the inhibitors of a VEGF receptor, as disclosed herein, containing only the extracellular region or containing, in addition, the transmembrane region may be constructed which have substantially the same activity.

The cloned sVEGF-R cDNA obtained through the methods described above may be recombinantly expressed by molecular cloning into an expression vector containing a suitable promoter and other appropriate transcription regulatory elements, and transferred into prokaryotic or eukaryotic host cells to produce recombinant sVEGF-R. Techniques for such manipulations are fully described in Maniatis, T. *et al.*, *supra*, and are well known in the art.

Expression vectors are defined herein as DNA sequences that are required for the transcription of cloned copies of genes and the translation of their mRNAs in an appropriate host. Such vectors can be used to express eukaryotic genes in a variety of hosts such as bacteria, bluegreen algae, fungal cells, yeast cells, plant cells, insect cells and animal cells.

Specifically designed vectors allow the shuttling of DNA between hosts such as bacteria-yeast

- 13 -

or bacteria-animal or bacteria-insect cells. An appropriately constructed expression vector should contain: an origin of replication for autonomous
5 replication in host cells, selectable markers, a limited number of useful restriction enzyme sites, a potential for high copy number, and active promoters. A promoter is defined as a DNA sequence that directs RNA polymerase to bind to DNA and initiate RNA
10 synthesis. A strong promoter is one which causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors, specifically designed plasmids or viruses.

15 A variety of mammalian expression vectors may be used to express recombinant sVEGF-R in mammalian cells. Commercially available mammalian expression vectors which may be suitable for recombinant sVEGF-R expression, include but are not limited to, pMCIneo
20 (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), EBO-pSV2-neo (ATCC 37593) pBPV-1(8-2) (ATCC 37110), pdBPV-MMTneo(342-12) (ATCC 37224), pRSVgpt (ATCC 37199), pRSVneo (ATCC 37198), pSV2-dhfr (ATCC 37146), pUCTag (ATCC 37460), and gZD35 (ATCC 37565).

25 DNA encoding sVEGF-R may also be cloned into an expression vector for expression in a recombinant host cell. Recombinant host cells may be prokaryotic or eukaryotic, including but not limited to bacteria, yeast, mammalian cells including but not limited to
30 cell lines of human, bovine, porcine, monkey and rodent origin, and insect cells including but not limited to

- 14 -

drosophila, moth, mosquito and armyworm derived cell lines. Cell lines derived from mammalian species which may be suitable and which are commercially available, include but are not limited to, CV-1 (ATCC CCL 70), COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCC CCL 2), C1271 (ATCC CRL 1616), BS-C-1 (ATCC CCL 26) and MRC-5 (ATCC CCL 171). Insect cell lines which may be suitable and are commercially available include but are not limited to 3M-S (ATCC CRL 8851) moth (ATCC CCL 80) mosquito (ATCC CCL 194 and 195; ATCC CRL 1660 and 1591) and armyworm (Sf9, ATCC CRL 1711).

The expression vector may be introduced into host cells via any one of a number of techniques including but not limited to transformation, transfection, liposome or protoplast fusion, and electroporation. The expression vector-containing cells are clonally propagated and individually analyzed to determine whether they produce sVEGF-R protein. Identification of sVEGF-R expressing host cell clones may be done by several means, including but not limited to immunological reactivity with anti-sVEGF-R antibodies, binding to radiolabelled VEGF, and the presence of host cell-secreted sVEGF-R activity.

Expression of sVEGF-R DNA may also be performed using *in vitro* produced synthetic mRNA. Synthetic mRNA can be efficiently translated in various cell-free systems, including but not limited to wheat germ extracts and reticulocyte extracts, as well as

- 15 -

efficiently translated in cell based systems, including but not limited to microinjection into frog oocytes, with microinjection into frog oocytes being preferred.

5 Levels of sVEGF-R protein produced by host cells may be quantitated by immunoaffinity and/or ligand affinity techniques. sVEGF-R-specific affinity beads or sVEGF-R-specific antibodies are used to isolate ³⁵S-methionine labelled or unlabelled sVEGF-R
10 protein. Labelled sVEGF-R protein is analyzed by SDS-PAGE. Unlabelled sVEGF-R protein is detected by Western blotting, ELISA or RIA assays employing sVEGF-R specific antibodies, or by ligand blotting with labelled VEGF.

15 Following expression of sVEGF-R in a recombinant host cell, sVEGF-R protein may be recovered to provide sVEGF-R in active form, capable of binding VEGF without stimulating mitogenesis. Several sVEGF-R purification procedures are available and suitable for
20 use. sVEGF-R may be purified from cell lysates and extracts, or from conditioned culture medium, by various combinations of, or individual application of salt fractionation, ion exchange chromatography, size exclusion chromatography, hydroxylapatite adsorption
25 chromatography, reversed phase chromatography, heparin sepharose chromatography, VEGF ligand affinity chromatography, and hydrophobic interaction chromatography.

 In addition, recombinant sVEGF-R can be
30 separated from other cellular proteins by use of an immuno-affinity column made with monoclonal or

- 16 -

polyclonal antibodies specific for full length sVEGF-R, or polypeptide fragments of sVEGF-R.

- 5 Identification of sVEGF-RI - In an attempt to clone the VEGF receptor cDNA (flt) a HUVEC λ gt10 cDNA library was screened with a DNA probe derived from the extracellular domain of the membrane bound or full length form of this receptor as shown in Figure 1.
- 10 Four incomplete clones, all lacking various lengths of 5' coding sequence, were isolated from screening a total of 1×10^6 plaques. Two of these isolates represent partial clones that were identical to full length flt, one of which contained the complete 3'
- 15 coding region of the form described by Shibuya *et al.*, supra. The other two clones were identical to full length flt up to base pair number 2219 (Table 1 and Figure 2) where they then diverged from full length flt. These clones (clone 7 and 11) coded for an
- 20 additional unique 31 amino acids before the open reading frame is terminated by a TAA codon (Table 2 and Figure 3).

Clone 7 and 11 coded for a protein with a predicted molecular mass of about 75 kDa containing 12

25 putative N-linked glycosylation sites. This version of the receptor was missing the transmembrane and intracellular kinase domains and thus coded for a natural soluble form of the VEGF receptor (sVEGF-RI). Further, the protein molecule predicted by sVEGF-RI has

30 only the first six Ig-like domains, missing the one closest to the transmembrane sequence (Figure 1). The

- 17 -

31 amino acids at the C-terminal end of sVEGF-RI contain two cysteine residues, but does not resemble an Ig domain.

5

Expression of sVEGF-RI in Sf9 cells - To analyze the binding and biological properties of this form of the receptor, the protein was expressed using a baculovirus expression system. Clone 7 was missing about 350 base pairs of coding sequence at the 5' end. This region was cloned by PCR using the primers described above and in Example 1. A clone containing the complete coding region of sVEGF-RI was constructed by combining the 5' PCR fragment with sVEGF-RI clone 7 which overlapped at a SacI site. The 5' EcoRI site was then changed to a BamHI site and the full length sVEGF-RI was cloned into pBluebac III (Invitrogen) as a BamHI/BamHI fragment. A recombinant baculovirus P-3 stock containing the sVEGF-RI gene 3' in relation to the polyhedrin promoter was then prepared as described herein.

Culture media from small scale infections were tested for the ability to form high molecular weight complexes with [¹²⁵I]VEGF. The labeled ligand and culture media from the baculovirus infected cells were combined and incubated. The reactions were then analyzed by size exclusion chromatography. When the wild-type infected culture medium was mixed with the radioactive ligand (Figure 4) a single radioactive peak was observed. However, when the sVEGF-RI infected culture medium was used, a high molecular weight complex was formed, as evident by the appearance of a

- 18 -

second peak in this reaction eluting near the void volume of the column. This experiment showed that the natural soluble form of the FLT VEGF receptor, sVEGF-RI, forms a high molecular weight complex with VEGF.

The recombinantly produced sVEGF-R is purified from the recombinant host cell extracts or cell culture fluid using heparin-sepharose column chromatography which specifically binds the sVEGF-R protein. The heparin-sepharose bound VEGF-R column is washed using a suitable buffer containing between 0.1M and 0.6M NaCl which removes contaminating proteins without significant loss of sVEGF-R. The sVEGF-R is eluted from the heparin-sepharose column using a suitable buffer containing about 1M NaCl, yielding substantially purified sVEGF-R.

Binding of the sVEGF-RI to VEGF - The binding of ^{125}I -labelled VEGF to sVEGF-RI was characterized by crosslinking, and by complex formation with sVEGF-RI absorbed to 96 well plates.

The crosslinked products are shown in Figure 6. The sVEGF-RI was cross-linked to [^{125}I]VEGF (lane 1); in the presence of unlabelled VEGF (lane 2) and unlabelled bFGF (lane 3). Two high molecular weight bands (about 145 kDa and 245 kDa) were formed in the sVEGF-RI and [^{125}I]VEGF containing reaction, and in the sVEGF-RI and [^{125}I]VEGF plus an excess of unlabelled bFGF reaction. The two high molecular weight bands were not present when sVEGF-RI was

- 19 -

incubated with [125 I]VEGF plus an excess of unlabelled VEGF, demonstrating the specificity of sVEGF-RI for VEGF, and the ability of sVEGF-RI to form a dimer. The
5 145 kDa band is presumably a crosslinked complex containing one receptor molecule (about 100 kDa) and a VEGF dimer (about 46 kDa). As shown in Figure 6 complexes containing two receptor molecules (about 245
10 kDa) were also observed. This suggests that each VEGF dimer can bind one or two receptor molecules and that the soluble form of the VEGF receptor may undergo ligand-induced dimerization.

The affinity of sVEGF-RI for VEGF was evaluated by absorbing sVEGF-RI to the surface of a 96
15 well plate, followed by blocking the nonspecific sites with 0.5% gelatin. Variable amounts of labeled ligand were added to each well. These results demonstrate that sVEGF-RI binds VEGF with high affinity with an apparent K_d of about 20pM (Figure 7). Since the
20 soluble form of the receptor is missing the Ig domain closest to the transmembrane spanning region, this domain is not required for ligand binding.

The sVEGF-RI is shown to inhibit binding of VEGF to HUVECs by incubating cultured HUVECs with
25 [125 I]VEGF and various amounts of sVEGF-RI. Following incubation, the cells are washed to remove unbound [125 I]VEGF. The cells are then solubilized and the amount of cell-associated 125 I is determined by gamma
30 counter, which demonstrates the amount of [125 I]VEGF which was capable of binding to the cellular VEGF receptor in the presence of sVEGF-RI. Using this

- 20 -

method, it is demonstrated that sVEGF-RI was capable of inhibiting [125 I]VEGF binding to HUVECs VEGF receptor (see Figure 8).

5 Since sVEGF-RI was able to inhibit VEGF binding to cell receptors, it was then determined that sVEGF-RI could inhibit VEGF induced mitogenesis. Cells are preincubated with sVEGF-RI and then incubated with VEGF in the presence of [3 H]thymidine. Following
10 incubation, the amount of cellular DNA-incorporated [3 H]thymidine is measured which indicates whether VEGF has induced mitogenesis and caused [3 H]thymidine to be incorporated into cellular DNA. The presence of sVEGF-RI inhibits the ability of VEGF to stimulate
15 mitogenesis as shown in Figure 9.

 The inhibitor of the present invention can be used for the inhibition of VEGF activity. The inhibitor can be used either topically or intravascularly. For topical applications the
20 formulation would be applied directly at a rate of about 10 ng to about 1 mg/cm²/day. For intravaneous applications, the inhibitor is used at a rate of about 1 µg to about 10 mg/kg/day of body weight. For
25 internal use, the formulation may be released directly into the region to be treated either from implanted slow release polymeric material or from slow release pumps or repeated injections. The release rate in either case is about 100 ng to about 100 µg/day/cm³.

 For non-topical application the VEGF
30 inhibitor is administered in combination with pharmaceutically acceptable carriers or diluents such

- 21 -

as phosphate buffer, saline, phosphate buffered saline, Ringer's solution, and the like, in a pharmaceutical composition, according to standard pharmaceutical practice. For topical application, various pharmaceutical formulations are useful for the administration of the active compound of this invention. Such formulations include, but are not limited to, the following: ointments such as hydrophilic petrolatum or polyethylene glycol ointment; pastes which may contain gums such as xanthan gum; solutions such as alcoholic or aqueous solutions; gels such as aluminum hydroxide or sodium alginate gels; albumins such as human or animal albumins; collagens such as human or animal collagens; celluloses such as alkyl celluloses, hydroxy alkyl celluloses and alkylhydroxyalkyl celluloses, for example methylcellulose, hydroxyethyl cellulose, carboxymethyl cellulose, hydroxypropyl methylcellulose, and hydroxypropyl cellulose; polyoxamers such as Pluronic® Polyols exemplified by Pluronic® F-127; tetronics such as tetronic 1508; and alginates such as sodium alginate.

The following examples are provided as illustrative of the present invention without, however, limiting the same thereto.

EXAMPLE 1

Cloning flt-related sVEGF-RI - A 580 base pair DNA probe for flt was obtained by PCR of the HUVEC phage library using the primers 5' GCACCTTGTTGTGGCTGAC 3'

- 22 -

(SEQ. ID. No.: 1) and 5' TGGAATTCGTGCTGCTTCCTGGTCC
3' (SEQ. ID. No.: 2). The resulting DNA fragment was
cloned into pGEM3Z as a XbaI/EcoRI fragment. The
5 probe was prepared by the random priming method
[Feinberg, A.P. and Vogelstein, B., (1983)
Anal.Biochem., 132, pp.6-13] using the megaprime kit
(Amersham) at a specific activity of 1×10^7 cpm/ng.
The HUVEC cDNA library was plated at a density of $5 \times$
10 10^4 plaques/150 cm plate then about 1×10^6 plaques
were screened by hybridization as previously described
[Maniatis, T. et al., supra]. Briefly, following
prehybridization at 42°C for 2 hours in 50% formamide,
5X SSC, 5X Denhardt's solution, 0.1% SDS, 100 µg/ml
15 salmon sperm DNA (hybridization buffer) the filters
were hybridized with the probe for 16 hours at 42°C in
hybridization buffer. The filters were washed one
time for 15 min at room temperature in 2X SSC then
three times at 55°C in 0.1 X SSC. Four positive
20 plaques were identified and rescreened two additional
times to obtain homogeneous isolates. Inserts were
cloned into pGEM3Z for DNA sequence analysis. Two of
these clones were identified which contained less than
the full length flt coding region. DNA sequence
25 analysis showed that these clones lacked the 5' coding
region of flt. The DNA sequence is shown in Table 1
and Figure 2, and the deduced amino acid sequence is
shown in Table 2 and Figure 3. The 5' end of flt was
cloned by PCR using the primers 5'
30 GGAATTCGCGCTCACCATGGTCAGC 3' (SEQ.ID.NO.:3) and 5'
TTTGAATTCACCCGGCAGGGAATGACG 3' (SEQ.ID.NO.:4). The
PCR fragment generated with this set of primers was
cloned into flt clone 7 as an EcoRI/SacI fragment.

- 23 -

TABLE 1

5 GCGGACACTCTCTCGGCTCCTCCCCGGCAGCGGCGGGCTCGGAGCGGGCTCCGGGG
CTCGGGTGCAGCGGCCAGCGGGCCTGCGGGCGAGGATTACCGGGGAAGTGGTTGTCTC
CTGGCTGGAGCCGCCAGACGGCCCTCAGGGCGGGGGCCGGCGGGCGGAACGAGAGG
10 ACGGACTCTGGCGGCCGGGTGCTTGGCCGGGGAGCGCGGCACCGGGCGAGCAGGCCG
CGTCGCGCTCACC ATG GTC AGC TAC TGG GAC ACC GGG GTC CTG CTG
TGC GCG CTG CTC AGC TGT CTG CTT CTC ACA GGA TCT AGT TCA GGT
15 TCA AAA TTA AAA GAT CCT GAA CTG AGT TTA AAA GGC ACC CAG CAC
ATC ATG CAA GCA GGC CAG ACA CTG CAT CTC CAA TGC AGG GGG GAA
20 GCA GCC CAT AAA TGG TCT TTG CCT GAA ATG CTG AGT AAG GAA ACC
GAA AGG CTG AGC ATA ACT AAA TCT GCC TGT GGA AGA AAT GGC AAA

25

30

- 24 -

CAA TTC TGC AGT ACT TTA ACC TTG AAC ACA GCT CAA GCA AAC CAC
ACT GGC TTC TAC AGC TGC AAA TAT CTA GCT GTA CCT ACT TCA AAG
5 AAG AAG GAA ACA GAA TCT GCA ATC TAT ATA TTT ATT AGT GAT ACA
GGT AGA CCT TTC GTA GAG ATG TAC AGT GAA ATC CCC GAA ATT ATA
10 CAC ATG ACT GAA GGA AGG GAG CTC GTC ATT CCC TGC CGG GTT ACG
TCA CCT AAC ATC ACT GTT ACT TTA AAA AAG TTT CCA CTT GAC ACT
TTG ATC CCT GAT GGA AAA CGC ATA ATC TGG GAC AGT AGA AAG GGC
15 TTC ATC ATA TCA AAT GCA ACG TAC AAA GAA ATA GGG CTT CTG ACC
TGT GAA GCA ACA GTC AAT GGG CAT TTG TAT AAG ACA AAC TAT CTC
20 ACA CAT CGA CAA ACC AAT ACA ATC ATA GAT GTC CAA ATA AGC ACA

25

30

- 25 -

CCA CGC CCA GTC AAA TTA CTT AGA GGC CAT ACT CTT GTC CTC AAT
TGT ACT GCT ACC ACT CCC TTG AAC ACG AGA GTT CAA ATG ACC TGG
5 AGT TAC CCT GAT GAA AAA AAT AAG AGA GCT TCC GTA AGG CGA CGA
ATT GAC CAA AGC AAT TCC CAT GCC AAC ATA TTC TAC AGT GTT CTT
10 ACT ATT GAC AAA ATG CAG AAC AAA GAC AAA GGA CTT TAT ACT TGT
CGT GTA AGG AGT GGA CCA TCA TTC AAA TCT GTT AAC ACC TCA GTG
CAT ATA TAT GAT AAA GCA TTC ATC ACT GTG AAA CAT CGA AAA CAG
15 CAG GTG CTT GAA ACC GTA GCT GGC AAG CGG TCT TAC CGG CTC TCT
ATG AAA GTG AAG GCA TTT CCC TCG CCG GAA GTT GTA TGG TTA AAA
20 GAT GGG TTA CCT GCG ACT GAG AAA TCT GCT CGC TAT TTG ACT CGT

25

30

- 26 -

GGC TAC TCG TTA ATT ATC AAG GAC GTA ACT GAA GAG GAT GCA GGG
AAT TAT ACA ATC TTG CTG AGC ATA AAA CAG TCA AAT GTG TTT AAA
5 AAC CTC ACT GCC ACT CTA ATT GTC AAT GTG AAA CCC CAG ATT TAC
GAA AAG GCC GTG TCA TCG TTT CCA GAC CCG GCT CTC TAC CCA CTG
10 GGC AGC AGA CAA ATC CTG ACT TGT ACC GCA TAT GGT ATC CCT CAA
CCT ACA ATC AAG TGG TTC TGG CAC CCC TGT AAC CAT AAT CAT TCC
GAA GCA AGG TGT GAC TTT TGT TCC AAT AAT GAA GAG TCC TTT ATC
15 CTG GAT GCT GAC AGC AAC ATG GGA AAC AGA ATT GAG AGC ATC ACT
CAG CGC ATG GCA ATA ATA GAA GGA AAG AAT AAG ATG GCT AGC ACC
20 TTG GTT GTG GCT GAC TCT AGA ATT TCT GGA ATC TAC ATT TGC ATA

25

30

- 27 -

GCT TCC AAT AAA GTT GGG ACT GTG GGA AGA AAC ATA AGC TTT TAT
ATC ACA GAT GTG CCA AAT GGG TTT CAT GTT AAC TTG GAA AAA ATG
5 CCG ACG GAA GGA GAG GAC CTG AAA CTG TCT TGC ACA GTT AAC AAG
TTC TTA TAC AGA GAC GTT ACT TGG ATT TTA CTG CGG ACA GTT AAT
10 AAC AGA ACA ATG CAC TAC AGT ATT AGC AAG CAA AAA ATG GCC ATC
ACT AAG GAG CAC TCC ATC ACT CTT AAT CTT ACC ATC ATG AAT GTT
TCC CTG CAA GAT TCA GGC ACC TAT GCC TGC AGA GCC AGG AAT GTA
15 TAC ACA GGG GAA GAA ATC CTC CAG AAG AAA GAA ATT ACA ATC AGA
GGT GAG CAC TGC AAC AAA AAG GCT GTT TTC TCT CGG ATC TCC AAA
20 TTT AAA AGC ACA AGG AAT GAT TGT ACC ACA CAA AGT AAT GTA AAA
CAT TAA

25

30

- 28 -

AGGACTCATTAAAAAGTAACAGTTGTCTCATATCATCTTGATTTATTGTCACTGTTG

CTAACTTTCAGGCTCGGAGGAGATGCTCCTCCCAAATGAGTTCGGAGATGATAGCA

5

GTAATAATGAGACCCCGGGCTCCAGCTCTGGGCCCCCATTGAGCCGAGGGGGCT

GCTCCGGGGGGCGACTTGCTGCACGTTTGGATTTGGAGGATCCCTGCACTGCCTTC

10 TCTGTGTTTGTGCTCTTGCTGTTTTCTCCTGCCTGATAACAACAACCTGGGATGA

TCCTTTCATTTTGATGCCAACCTCTTTTATTTTAAAGCGGGCCCTATAGT

(SEQ. ID. NO.: 5)

15

20

25

30

- 29 -

TABLE 2

5 Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu
Cys Ala Leu Leu Ser Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly
Ser Lys Leu Lys Asp Pro Glu Leu Ser Leu Lys Gly Thr Gln His
10 Ile Met Gln Ala Gly Gln Thr Leu His Leu Gln Cys Arg Gly Glu
Ala Ala His Lys Trp Ser Leu Pro Glu Met Val Ser Lys Glu Ser
Glu Arg Leu Ser Ile Thr Lys Ser Ala Cys Gly Arg Asn Gly Lys
15 Gln Phe Cys Ser Thr Leu Thr Leu Asn Thr Ala Gln Ala Asn His
Thr Gly Phe Tyr Ser Cys Lys Tyr Leu Ala Val Pro Thr Ser Lys
20 Lys Lys Glu Thr Glu Ser Ala Ile Tyr Ile Phe Ile Ser Asp Thr
Gly Arg Pro Phe Val Glu Met Tyr Ser Glu Ile Pro Glu Ile Ile

25

30

- 30 -

His Met Thr Glu Gly Arg Glu Leu Val Ile Pro Cys Arg Val Thr
Ser Pro Asn Ile Thr Val Thr Leu Lys Lys Phe Pro Leu Asp Thr
5 Leu Ile Pro Asp Gly Lys Arg Ile Ile Trp Asp Ser Arg Lys Gly
Phe Ile Ile Ser Asn Ala Thr Tyr Lys Glu Ile Gly Leu Leu Thr
10 Cys Glu Ala Thr Val Asn Gly His Leu Tyr Lys Thr Asn Tyr Leu
Thr His Arg Gln Thr Asn Thr Ile Ile Asp Val Gln Ile Ser Thr
Pro Arg Pro Val Lys Leu Leu Arg Gly His Thr Leu Val Leu Asn
15 Cys Thr Ala Thr Thr Pro Leu Asn Thr Arg Val Gln Met Thr Trp
Ser Tyr Pro Asp Glu Lys Asn Lys Arg Ala Ser Val Arg Arg Arg
20 Ile Asp Gln Ser Asn Ser His Ala Asn Ile Phe Tyr Ser Val Leu

25

30

- 31 -

Thr Ile Asp Lys Met Gln Asn Lys Asp Lys Gly Leu Tyr Thr Cys
Arg Val Arg Ser Gly Pro Ser Phe Lys Ser Val Asn Thr Ser Val
5 His Ile Tyr Asp Lys Ala Phe Ile Thr Val Lys His Arg Lys Gln
Gln Val Leu Glu Thr Val Ala Gly Lys Arg Ser Tyr Arg Leu Ser
10 Met Lys Val Lys Ala Phe Pro Ser Pro Glu Val Val Trp Leu Lys
Asp Gly Leu Pro Ala Thr Glu Lys Ser Ala Arg Tyr Leu Thr Arg
Gly Tyr Ser Leu Ile Ile Lys Asp Val Thr Glu Glu Asp Ala Gly
15 Asn Tyr Thr Ile Leu Leu Ser Ile Lys Gln Ser Asn Val Phe Lys
Asn Leu Thr Ala Thr Leu Ile Val Asn Val Lys Pro Gln Ile Tyr
20 Glu Lys Ala Val Ser Ser Phe Pro Asp Pro Ala Leu Tyr Pro Leu

25

30

- 32 -

Gly Ser Arg Gln Ile Leu Thr Cys Thr Ala Tyr Gly Ile Pro Gln
Pro Thr Ile Lys Trp Phe Trp His Pro Cys Asn His Asn His Ser
5 Glu Ala Arg Cys Asp Phe Cys Ser Asn Asn Glu Glu Ser Phe Ile
Leu Asp Ala Asp Ser Asn Met Gly Asn Arg Ile Glu Ser Ile Thr
10 Gln Arg Met Ala Ile Ile Glu Gly Lys Asn Lys Met Ala Ser Thr
Leu Val Val Ala Asp Ser Arg Ile Ser Gly Ile Tyr Ile Cys Ile
Ala Ser Asn Lys Val Gly Thr Val Gly Arg Asn Ile Ser Phe Tyr
15 Ile Thr Asp Val Pro Asn Gly Phe His Val Asn Leu Glu Lys Met
Pro Thr Glu Gly Glu Asp Leu Lys Leu Ser Cys Thr Val Asn Lys
20 Phe Leu Tyr Arg Asp Val Thr Trp Ile Leu Leu Arg Thr Val Asn

25

30

- 33 -

Asn Arg Thr Met His Tyr Ser Ile Ser Lys Gln Lys Met Ala Ile
Thr Lys Glu His Ser Ile Thr Leu Asn Leu Thr Ile Met Asn Val
5 Ser Leu Gln Asp Ser Gly Thr Tyr Ala Cys Arg Ala Arg Asn Val
Tyr Thr Gly Glu Glu Ile Leu Gln Lys Lys Glu Ile Thr Ile Arg
10 Gly Glu His Cys Asn Lys Lys Ala Val Phe Ser Arg Ile Ser Lys
Phe Lys Ser Thr Arg Asn Asp Cys Thr Thr Gln Ser Asn Val Lys
His *** (SEQ. ID. NO.: 6)
15

EXAMPLE 2

Expression of sVEGF-RI in Sf9 insect cells - The full
20 length sequence encoding sVEGF-RI was cloned as an
EcoRI/BamHI fragment into pGEM3Z. The EcoRI site was
then modified to a BamHI site and cloned into pBlueBac
III 3' of the polyhedrin promoter (psFLTblue). This
25 plasmid was transfected into Sf9 armyworm cells using
liposomes. After 48 hours the medium from the
transfected cells which contains recombinant polyhedrin
virus particles, was harvested. Dilutions (10^3 - 10^4
fold) of the virus were prepared and plaque purified in
30 soft agar containing 150 µg/ml 5-bromo-4-chloro-3-

- 34 -

indolyl- β -D-galactoside. Recombinant plaques were identified by blue color and used to infect Sf9 cells (5×10^5 cells/well) in 12 well plates. Medium (100 5 μ l) from polyhedrin minus infections was used to prepare P-2 viral stocks by infecting 2.5×10^6 cells in a T-25 flask. Large scale high titer P-3 viral stocks were then prepared by infecting Sf9 cells (500 ml at 2×10^6 cells/ml) with 5 ml of the P-2 stock then 10 incubating at 27°C for 5 - 6 days and the medium was harvested by centrifugation. Protein expression was accomplished by infecting cells at a density of $2-2.5 \times 10^6$ cells/ml with a multiplicity of infection of 5 - 10. Twenty four hours after infection the cells were 15 changed to a serum free medium (SF900II, Gibco BRL), incubated for an additional 48 hours and the medium was collected. This conditioned medium contains the recombinantly expressed sVEGF-RI protein.

20

EXAMPLE 3

Iodination of VEGF - ^{125}I -labeled human recombinant VEGF was prepared by the chloramine T method (Hunter, W.M. and Greenwood, F.C., (1962) Nature (London), 194, 25 pp. 495-496). Briefly, 1 μ g of VEGF in 30% acetonitrile/0.1% trifluoroacetic acid was adjusted to pH 7.1 by the addition of 1/3 volume of 0.4 M sodium phosphate buffer, pH 7.1. Freshly dissolved chloramine T (4 μ l of a 2 mg/ml stock in 0.1 M sodium phosphate 30 buffer, pH 7.1) was added to the VEGF solution and reacted for 45 seconds at room temperature (total

- 35 -

volume of 150 μ l). The reaction was stopped by the addition of 50 μ l of 10 mM KI and 50 μ l of 2 mg/ml meta bisufite. The labeled ligand was separated from the free ^{125}I by gel filtration on a 0.7 X 15 cm Sephadex G-25 column equilibrated in PBS with 1 mg/ml gelatin. Fractions were counted in a Packard γ counter, aliquoted and stored at -70°C . VEGF was labeled to a specific activity of 5×10^5 to 1×10^6 cpm/ng.

Gel Filtration Chromatography - Receptor-ligand complex was formed by incubating 10 μ l of ^{125}I -labeled VEGF (10^5 cpm) with 100 μ l of either wild-type or baculovirus sVEGF-RI-containing, infected Sf9 cell culture medium overnight at room temperature. The reaction products were separated on a Sephacryl S200 gel filtration column (0.7 X 25 cm) equilibrated in PBS, 1 mg/ml gelatin, at a flow rate of 15 ml/hr. Fractions (0.75 ml) were collected and analyzed in a γ counter. Receptor-ligand complexes pass quickly through the column while the free labelled VEGF passes through more slowly. The results of this experiment shown in Figure 4 demonstrate the formation of a high molecular weight complex between labelled VEGF and sVEGF-RI protein. This shows that sVEGF-RI binds VEGF.

Crosslinking - Purified sVEGF-RI (1-10ng) was added to 25 μ l of binding buffer (Dulbecco's Modified Eagle's medium (DME), 25 mM HEPES, pH 7.5, 0.3% gelatin), and 1×10^5 cpm of [^{125}I]-VEGF was added (Figure 6, lane 1) with either 200ng of unlabelled VEGF (lane 2) or bFGF

- 36 -

(lane 3), then incubated 2 to 16 hours at room temperature. Bis(sulfosuccinimidyl)suberate (Pierce) crosslinker was added to a final concentration of 1 mM. The reaction was stopped after 15 min by the addition of boiling SDS PAGE sample buffer. The crosslinked products were separated by SDS PAGE on a 7.5% acrylamide gel and analyzed either by autoradiography or a phosphoimager. The results are shown in Figure 6 and demonstrate that sVEGF-RI binds labelled VEGF by the appearance of two bands of about 145 kDa and 245 kDa. The 145 kDa band consists of one sVEGF-RI molecule and one VEGF molecule (Monomer, M.). The 245 kDa band apparently consists of two sVEGF-RI molecules and one VEGF dimer (D). Free VEGF ligand (L) dimers migrated at about 45 kDa.

Binding assay - The binding of sVEGF-RI to VEGF was analyzed using a 96 well plate assay as described by Duan, D-S. R. *et al.*, supra. Briefly, sVEGF-RI, 50 to 200 μ l partially purified by Mono Q chromatography (Pharmacia), was diluted to 10 ml in 25 mM TRIS, pH 7.4, 100 mM NaCl, 20 mM NH_4HCO_3 . Aliquots (100 μ l) were absorbed to the surface of a 96 well plate for 18 hours at 4°C, the plates were then washed twice with blocking buffer (DME, 25 mM HEPES, pH 7.5, 0.5% gelatin) and the nonspecific sites were blocked in the same buffer for 6 hours at 4°C. The plate was then washed twice in binding buffer. Various amounts of [^{125}I]VEGF were added to the wells in a final volume of 100 μ l/well and incubated for 2 hours at room

- 37 -

temperature. The wells were washed three times with 100 μ l of binding buffer, the bound protein was solubilized with 100 μ l of 1% SDS, 0.5% BSA and counted in a γ counter. The results, shown in Figure 7, were analyzed by the method of Scatchard [Scatchard, G., (1949) Ann. N.Y. Acad. Sci., 51, pp. 660-672]. The analysis demonstrates that sVEGF-RI retains high affinity binding for VEGF with a K_d value of about 20 pM. This clearly demonstrates that sVEGF-RI, lacking the transmembrane region and adjacent Ig-like domain, binds VEGF with high affinity and that these regions are not required for VEGF binding.

15

EXAMPLE 4

Inhibition of VEGF binding by sVEGF-RI - The ability of sVEGF-RI to inhibit VEGF binding to HUVECs was tested. HUVECs were plated at 50,000 cells/well in 24 well plates precoated with gelatin, and allowed to grow to confluence. A constant amount of [125 I]VEGF (100,000 cpm) was mixed with various amounts of partially purified sVEGF-RI in binding buffer, in a total volume of 200 μ l and preincubated at room temperature for 1 hour. Samples were added to the cells and incubated for 4 hours at 4°C with shaking. The medium was then aspirated and the cells were washed three times with binding buffer. The bound radioactivity was solubilized with 50 mM TRIS-HCl, pH 8.0, 150 mM NaCl, 1% NP40, 1% BSA and counted in a γ counter.

- 38 -

The results are shown in Figure 8. At the highest concentration of sVEGF-RI, VEGF binding to HUVECs was reduced by 70%. It may, however, be difficult to completely inhibit binding to the cellular membrane bound receptor since one molecule of sVEGF-R bound to a VEGF dimer may be able to bind to cell associated receptor to form an inactive (sVEGF-RI)-VEGF-(membrane spanning VEGF receptor) complex.

EXAMPLE 5

Inhibition of VEGF mediated mitogenesis by sVEGF-RI
Mitogenic inhibition - Since sVEGF-RI was able to inhibit VEGF binding to endothelial cells, it was then determined that the soluble receptor could inhibit VEGF induced mitogenesis in HUVECs. HUVECs were plated in gelatin coated 96 well plates at a density of 4000 cells/well in 100 μ l of DME supplemented with 10% heat inactivated fetal calf serum plus antibiotics (penicillin G, 100 units/ml; streptomycin sulfate, 100 μ g/ml). After 16 hours the medium was changed and test samples were added, cells were preincubated with a variable amount of purified sVEGF-RI for 15 minutes at 37°C before growth factor (10 ng/ml) was added. The cells were incubated for 24 hours then [methyl-³H]thymidine (0.8 μ Ci/well; 20 Ci/mmol: 1Ci = 37 GBq, final specific activity of 0.8 μ Ci/nmole) was added followed by incubated for an additional 72 hours at 37°C under 5% CO₂. The cells were then washed twice with Hank's balanced salt solution adjusted to pH 7.5

- 39 -

with 25 mM Hepes, 0.1% BSA. The cells were then lysed, the DNA was solubilized with 0.2 M Na_2CO_3 , 0.1 M NaOH, and [^3H]thymidine incorporation was quantified by scintillation counting. The results are shown in Figure 9. sVEGF-RI was able to completely inhibit VEGF induced [^3H]thymidine incorporation in HUVECs.

EXAMPLE 6

Purification of baculovirus expressed sVEGF-RI from Sf9 cells - Culture medium from Sf9 cells infected with a baculovirus construct designed to express sVEGF-RI (Example 2) was chromatographed through a heparin Sepharose CL-6B (Pharmacia) column (0.7 X 4 cm). The column was washed with 5 volumes of 10 mM Na-phosphate buffer, pH 6.2, 0.1 M NaCl, followed by 6 ml of 10 mM Na-phosphate buffer, pH 6.2, 0.6 M NaCl. The sVEGF-RI was eluted with 10 mM Na-phosphate buffer, pH 6.2, 1.0 M NaCl. Polyacrylamide gel electrophoresis was performed which demonstrated greater than 90% purity (as judged by coomassie blue staining) of the recombinantly produced sVEGF-R (Figure 5). The identity of the protein was confirmed by N-terminal protein sequence analysis. The actual N-terminus (Ser Lys Leu ...) of the recombinant protein differs by two amino acids from that predicted by Shibuya et al., supra. (Ser-Ser-Ser...). The peptidase cleavage site in sVEGF-RI produced in Sf9 cells was between residues gly-26 and ser-27.

- 40 -

EXAMPLE 7

Construction of KDR-related sVEGF-R - Soluble forms of
5 KDR (a known VEGF receptor) [Terman, B.I. et al.,
(1991) Oncogene 6, pp. 1677-1683; Terman, B.I. et al.,
(1992) Biochem. Biophys. Res. Comm. 187, pp. 1579-1586]
may exist naturally but have not yet been identified.
A soluble form of KDR is recombinantly constructed by
10 modifying its coding sequence by PCR using the primers
1) 5' TTTTGGATCCCTGCAGACAGATCTACGTTTGAGAACC 3' (SEQ.
ID. NO.: 7) and 2) 5' TTTTGGATCCTTAACGCTCTAGGACTGTGAGC
3' (SEQ. ID. NO.: 8), and pKDRA (the XhoI/EcoRI
15 fragment coding for the extracellular and transmembrane
domain of KDR cloned into the EcoRI site of pGEM 7Z
obtained from Promega) as a template (Figure 17). This
generated a translation stop codon after amino acid
residue number 663 of KDR which corresponds to the
extracellular domain of full length KDR. This modified
20 fragment is then used to replace the PstI/BamHI
fragment of pKDRA generating a truncated form of the
KDR gene (Figure 10) which codes for a soluble receptor
denoted sVEGF-RII (Figure 11). The XhoI site at base
pair number 257 is then changed to a BamHI site by
25 standard cloning techniques. Another truncated form of
the KDR receptor is created with primer 1 shown above,
and primer 3) 5' TTTTGGATCCAACGGTCCCTAGGATGATGAC 3' ,
(SEQ. ID. NO.: 9) (Figure 12). This form of KDR,
denoted sVEGF-RTMII, is truncated at the C-terminal
30 side of the transmembrane domain and therefore retains
the transmembrane region (Figure 13). A similar form
of the FLT receptor is generated by PCR using the

- 41 -

primers 4) 5' AGCACCTTGGTTGTGGCTGACTC 3' (SEQ. ID. NO.: 10) and 5) 5' TTTTGGATCCTTAGATAAGGAGGGTTAATAGG 3' (SEQ. ID. NO.: 11) and plasmid pmFLT (full length flt cloned into the EcoRI site of pGEM3Z obtained from Promega) as a template (Figure 16). The 780 base pair PCR fragment can then be cloned together with the EcoRI/XbaI fragment from pmFLT to produce an EcoRI/BAMHI fragment (Figure 14) encoding a truncated form of FLT (denoted sVEGF-RTMI) which retains the transmembrane domain but lacks the cytoplasmic domain (Figure 15). The EcoRI site at the 5' end of the gene is then modified to a BamHI site. The resulting truncated forms of KDR and FLT are then cloned into pBluebac111 (Stratagene) for expression in Sf9 insect cells. Characterization of these constructed truncated forms of VEGF receptors is accomplished by the techniques used to characterize sVEGF-RI as in Examples 2, 3, 4, 5, and 6.

20

25

30

- 42 -

SEQUENCE LISTING

5

(1) GENERAL INFORMATION:

(i) APPLICANT: Thomas, Kenneth A.
Kendall, Richard L.

10

(ii) TITLE OF INVENTION: INHIBITOR OF VASCULAR ENDOTHELIAL CELL
GROWTH FACTOR

15

(iii) NUMBER OF SEQUENCES: 18

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Merck & Co., Inc.
(B) STREET: P.O. Box 2000 126 E Lincoln Avenue
(C) CITY: Rahway
(D) STATE: NJ
(E) COUNTRY: USA
(F) ZIP: 07065-0907

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

25

30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

- 43 -

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Wallen, John W.III
(B) REGISTRATION NUMBER: 35,403
(C) REFERENCE/DOCKET NUMBER: 18888

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (908) 594-3905
(B) TELEFAX: (908) 594-4720

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCACCTTG6T TGTGGCTGAC

20

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 44 -

(ii) MOLECULE TYPE: cDNA

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TGGAAATTCGT GCTGCTTCCT GGTC

25

10

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

15

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGAATTCGCG GCTCACCATG GTCAGC

26

25

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

30

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

- 45 -

(ii) MOLECULE TYPE: cDNA

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TTTGAATTCA CCCGGCAGGG AATGACG

27

10

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2313 base pairs

(B) TYPE: nucleic acid

15

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

25

GCGGACACTC CTCTCGGCTC CTCCCCGGCA GCGGCGGCGG CTCGGAGCGG GCTCCGGGGC 60

TCGGGTGCAG CGGCCAGCGG GCCTGGCGGC GAGGATTACC CGGGGAAGTG GTTGTCTCCT 120

GGCTGGAGCC GCGAGACGGG CGCTCAGGCG GCGGGGCCGG CGGCGGCGAA CGAGAGGACG 180

30

GACTCTGGCG GCCGGGTCGT TGGCCGGGGG AGCGCGGGCA CCGGCGGAGC AGGCCGCGTC 240

GCGCTCACCA TGGTCAGCTA CTGGGACACC GGGGTCCTGC TGTGCGCGCT GCTCAGCTGT 300

- 46 -

	CTGCTTCTCA CAGGATCTAG TTCAGGTTC AATTAAAAG ATCCTGAACT GAGTTAAAA	360
5	GGCACCCAGC ACATCATGCA AGCAGGCCAG AACTGCATC TCCAATGCAG GGGGAAGCA	420
	GCCCATAAAT GGTCTTTGCC TGAATGGTG AGTAAGGAAA GCGAAAGGCT GAGCATACT	480
	AAATCTGCCT GTGGAAGAAA TGGCAACAA TTCTGCAGTA CTTAACCTT GAACACAGCT	540
10	CAAGCAAACC AACTGGCTT CTACAGETGC AAATATCTAG CTGTACCTAC TTCAAAGAAG	600
	AAGGAAACAG AATCTGCAAT CTATATATT ATTAGTGATA CAGGTAGACC TTTCGTAGAG	660
15	ATGTACAGTG AATCCCCGA AATTATACAC ATGACTGAAG GAAGGGAGCT CGTCATTCCC	720
	TGCCGGGTTA CGTCACCTAA CATCACTGT ACTTTAAAA AGTTTCCACT TGACACTTTG	780
	ATCCCTGATG GAAAACGCAT AATCTGGGAC AGTAGAAGG GCTTCATCAT ATCAAATGCA	840
20	ACGTACAAAG AAATAGGGCT TCTGACCTGT GAAGCAACAG TCAATGGGCA TTTGTATAAG	900
	ACAACTATC TCACACATCG ACAAACCAAT ACAATCATAG ATGTCCAAAT AAGCACACCA	960
25	CGCCAGTCA AATTACTTAG AGGCCATACT CTTGTCTCA ATTGTACTGC TACCACTCCC	1020
	TTGAACAGG GAGTTCAAAT GACCTGGAGT TACCCTGATG AAAAAATAA GAGAGCTTCC	1080
	GTAAGGCGAC GAATTGACCA AAGCAATTCC CATGCCAACA TATTCTACAG TGTCTTACT	1140
30	ATTGACAAAA TGCAGAACA AGACAAAGGA CTTTATACTT GTCGTGTAAG GAGTGGACCA	1200
	TCATTCAAAT CTGTTAACAC CTCAGTGCAT ATATATGATA AAGCATTCACT CACTGTGAAA	1260

- 47 -

	CATCGAAAAC AGCAGGTGCT TGA AACCGTA GCTGGCAAGC GGTCTTACC6 GCTCTCTATG	1320
5	AAAGTGAAGG CATTTCCTC GCCGGAAGTT STATGGTTAA AAGATGGGT ACCTGCGACT	1380
	GAGAAATCTG CTCGCTATTT GACTCGTGGC TACTCGTTAA TTATCAAGGA CGTAACTGAA	1440
	GAGGATGCAG GGAATTATAC AATCTTGCTG AGCATAAAAC AGTCAAATGT GTTTAAAAAC	1500
10	CTCACTGCCA CTCTAATTGT CAATGTGAAA CCCAGATTT ACGAAAAGGC CGTGTCATCG	1560
	TTCCAGACC CGGCTCTCTA CCCACTGGGC AGCAGACAAA TCCTGACTTG TACCGCATAT	1620
15	GGTATCCCTC AACCTACAAT CAAGTGGTTC TGGCACCCT GTAACCATAA TCATTCCGAA	1680
	GCAAGGTGTG ACTTTTGTTT CAATAATGAA GAGTCCTTTA TCCTGGATGC TGACAGCAAC	1740
	ATGGGAAACA GAATTGAGAG CATCACTCAG CGCATGGCAA TAATAGAAGG AAAGAATAAG	1800
20	ATGGCTAGCA CCTTGTTGT GGCTGACTCT AGAATTTCTG GAATCTACAT TTGCATAGCT	1860
	TCCAATAAAG TTGGGACTGT GGAAGAAAC ATAAGCTTTT ATATCACAGA TGTGCCAAAT	1920
25	GGGTTTCATG TTAAGTTGGA AAAATGCCG ACGGAAGGAG AGGACCTGAA ACTGTCTTGC	1980
	ACAGTTAACA AGTTCTTATA CAGAGACGTT ACTTGGATTT TACTGCGGAC AGTTAATAAC	2040
	AGAACAATGC ACTACAGTAT TAGCAAGCAA AAAATGGCCA TCACTAAGGA GCACTCCATC	2100
30	ACTCTTAATC TTACCATCAT GAATGTTTCC CTGCAAGATT CAGGCACCTA TGCCTGCAGA	2160
	GCCAGGAATG TATACACAGG GGAAGAAATC CTCCAGAAGA AAGAAATTAC AATCAGAGGT	2220

- 48 -

BAGCACTGCA ACAAAAAGGC TGTTTCTCT CGGATCTCCA AATTAAAAAG CACAAGGAAT 2280

GATTGTACCA CACAAAGTAA TGTAACAT TAA 2313

5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 687 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

20

Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu Cys Ala Leu Leu Ser
1 5 10 15

Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly Ser Lys Leu Lys Asp Pro
20 25 30

25

Glu Leu Ser Leu Lys Gly Thr Gln His Ile Met Gln Ala Gly Gln Thr
35 40 45

30

Leu His Leu Gln Cys Arg Gly Glu Ala Ala His Lys Trp Ser Leu Pro
50 55 60

Glu Met Val Ser Lys Glu Ser Glu Arg Leu Ser Ile Thr Lys Ser Ala
65 70 75 80

- 49 -

	Cys Gly Arg Asn Gly Lys Gln Phe Cys Ser Thr Leu Thr Leu Asn Thr	
	85	90 95
5	Ala Gln Ala Asn His Thr Gly Phe Tyr Ser Cys Lys Tyr Leu Ala Val	
	100	105 110
	Pro Thr Ser Lys Lys Lys Glu Thr Glu Ser Ala Ile Tyr Ile Phe Ile	
10	115	120 125
	Ser Asp Thr Gly Arg Pro Phe Val Glu Met Tyr Ser Glu Ile Pro Glu	
	130	135 140
15	Ile Ile His Met Thr Glu Gly Arg Glu Leu Val Ile Pro Cys Arg Val	
	145	150 155 160
	Thr Ser Pro Asn Ile Thr Val Thr Leu Lys Lys Phe Pro Leu Asp Thr	
	165	170 175
20	Leu Ile Pro Asp Gly Lys Arg Ile Ile Trp Asp Ser Arg Lys Gly Phe	
	180	185 190
	Ile Ile Ser Asn Ala Thr Tyr Lys Glu Ile Gly Leu Leu Thr Cys Glu	
25	195	200 205
	Ala Thr Val Asn Gly His Leu Tyr Lys Thr Asn Tyr Leu Thr His Arg	
	210	215 220
30	Gln Thr Asn Thr Ile Ile Asp Val Gln Ile Ser Thr Pro Arg Pro Val	
	225	230 235 240
	Lys Leu Leu Arg Gly His Thr Leu Val Leu Asn Cys Thr Ala Thr Thr	
	245	250 255

- 50 -

	Pro Leu Asn Thr Arg Val Gln Met Thr Trp Ser Tyr Pro Asp Glu Lys	
	260	265 270
5	Asn Lys Arg Ala Ser Val Arg Arg Arg Ile Asp Gln Ser Asn Ser His	
	275	280 285
	Ala Asn Ile Phe Tyr Ser Val Leu Thr Ile Asp Lys Met Gln Asn Lys	
10	290	295 300
	Asp Lys Gly Leu Tyr Thr Cys Arg Val Arg Ser Gly Pro Ser Phe Lys	
	305	310 315 320
	Ser Val Asn Thr Ser Val His Ile Tyr Asp Lys Ala Phe Ile Thr Val	
15	325	330 335
	Lys His Arg Lys Gln Gln Val Leu Glu Thr Val Ala Gly Lys Arg Ser	
	340	345 350
20	Tyr Arg Leu Ser Met Lys Val Lys Ala Phe Pro Ser Pro Glu Val Val	
	355	360 365
	Trp Leu Lys Asp Gly Leu Pro Ala Thr Glu Lys Ser Ala Arg Tyr Leu	
25	370	375 380
	Thr Arg Gly Tyr Ser Leu Ile Ile Lys Asp Val Thr Glu Glu Asp Ala	
	385	390 395 400
	Gly Asn Tyr Thr Ile Leu Leu Ser Ile Lys Gln Ser Asn Val Phe Lys	
30	405	410 415
	Asn Leu Thr Ala Thr Leu Ile Val Asn Val Lys Pro Gln Ile Tyr Glu	
	420	425 430

- 51 -

Lys Ala Val Ser Ser Phe Pro Asp Pro Ala Leu Tyr Pro Leu Gly Ser
435 440 445

5 Arg Gln Ile Leu Thr Cys Thr Ala Tyr Gly Ile Pro Gln Pro Thr Ile
450 455 460

Lys Trp Phe Trp His Pro Cys Asn His Asn His Ser Glu Ala Arg Cys
465 470 475 480

10 Asp Phe Cys Ser Asn Asn Glu Glu Ser Phe Ile Leu Asp Ala Asp Ser
485 490 495

Asn Met Gly Asn Arg Ile Glu Ser Ile Thr Gln Arg Met Ala Ile Ile
15 500 505 510

Glu Gly Lys Asn Lys Met Ala Ser Thr Leu Val Val Ala Asp Ser Arg
515 520 525

20 Ile Ser Gly Ile Tyr Ile Cys Ile Ala Ser Asn Lys Val Gly Thr Val
530 535 540

Gly Arg Asn Ile Ser Phe Tyr Ile Thr Asp Val Pro Asn Gly Phe His
25 545 550 555 560

Val Asn Leu Glu Lys Met Pro Thr Glu Gly Glu Asp Leu Lys Leu Ser
565 570 575

Cys Thr Val Asn Lys Phe Leu Tyr Arg Asp Val Thr Trp Ile Leu Leu
30 580 585 590

Arg Thr Val Asn Asn Arg Thr Met His Tyr Ser Ile Ser Lys Gln Lys
595 600 605

- 52 -

Met Ala Ile Thr Lys Glu His Ser Ile Thr Leu Asn Leu Thr Ile Met
610 615 620

5 Asn Val Ser Leu Gln Asp Ser Gly Thr Tyr Ala Cys Arg Ala Arg Asn
625 630 635 640

Val Tyr Thr Gly Glu Glu Ile Leu Gln Lys Lys Glu Ile Thr Ile Arg
645 650 655

10 Gly Glu His Cys Asn Lys Lys Ala Val Phe Ser Arg Ile Ser Lys Phe
660 665 670

Lys Ser Thr Arg Asn Asp Cys Thr Thr Gln Ser Asn Val Lys His
15 675 680 685

(2) INFORMATION FOR SEQ ID NO:7:

- 20 (1) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 25 (11) MOLECULE TYPE: DNA (genomic)

- 30 (x1) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TTTTGGATCC CTGCAGACAG ATCTACGTTT GAGAAC

- 53 -

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA (genomic)

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TTTTGGATCC TTAACGCTCT AGGACTGTGA GC

32

(2) INFORMATION FOR SEQ ID NO:9:

20 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TTTTGGATCC AACGGTCCCT AGGATGATGA C

31

- 54 -

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

5

(A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: DNA (genomic)

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:10:

15

AGCACCTTGG TTGTGGCTGA CTC

23

(2) INFORMATION FOR SEQ ID NO:11:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA (genomic)

30

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TTTGGATCC TTAGATAAGG AGGGTTAATA GG

32

- 55 -

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

5

(A) LENGTH: 661 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

15

Ser Lys Leu Lys Asp Pro Glu Leu Ser Leu Lys Gly Thr Gln His Ile

1

5

10

15

20

Met Gln Ala Gly Gln Thr Leu His Leu Gln Cys Arg Gly Glu Ala Ala

20

25

30

His Lys Trp Ser Leu Pro Glu Met Val Ser Lys Glu Ser Glu Arg Leu

35

40

45

25

Ser Ile Thr Lys Ser Ala Cys Gly Arg Asn Gly Lys Gln Phe Cys Ser

50

55

60

30

Thr Leu Thr Leu Asn Thr Ala Gln Ala Asn His Thr Gly Phe Tyr Ser

65

70

75

80

Cys Lys Tyr Leu Ala Val Pro Thr Ser Lys Lys Lys Glu Thr Glu Ser

85

90

95

- 56 -

	Ala Ile Tyr Ile Phe Ile Ser Asp Thr Gly Arg Pro Phe Val Glu Met	
	100	105 110
5	Tyr Ser Glu Ile Pro Glu Ile Ile His Met Thr Glu Gly Arg Glu Leu	
	115	120 125
	Val Ile Pro Cys Arg Val Thr Ser Pro Asn Ile Thr Val Thr Leu Lys	
	130	135 140
10	Lys Phe Pro Leu Asp Thr Leu Ile Pro Asp Gly Lys Arg Ile Ile Trp	
	145	150 155 160
	Asp Ser Arg Lys Gly Phe Ile Ile Ser Asn Ala Thr Tyr Lys Glu Ile	
15	165	170 175
	Gly Leu Leu Thr Cys Glu Ala Thr Val Asn Gly His Leu Tyr Lys Thr	
	180	185 190
20	Asn Tyr Leu Thr His Arg Gln Thr Asn Thr Ile Ile Asp Val Gln Ile	
	195	200 205
	Ser Thr Pro Arg Pro Val Lys Leu Leu Arg Gly His Thr Leu Val Leu	
	210	215 220
25	Asn Cys Thr Ala Thr Thr Pro Leu Asn Thr Arg Val Gln Met Thr Trp	
	225	230 235 240
	Ser Tyr Pro Asp Glu Lys Asn Lys Arg Ala Ser Val Arg Arg Arg Ile	
30	245	250 255
	Asp Gln Ser Asn Ser His Ala Asn Ile Phe Tyr Ser Val Leu Thr Ile	
	260	265 270

- 57 -

	Asp Lys Met Gln Asn Lys Asp Lys Gly Leu Tyr Thr Cys Arg Val Arg	
	275	280 285
5	Ser Gly Pro Ser Phe Lys Ser Val Asn Thr Ser Val His Ile Tyr Asp	
	290	295 300
	Lys Ala Phe Ile Thr Val Lys His Arg Lys Gln Gln Val Leu Glu Thr	
10	305	310 315 320
	Val Ala Gly Lys Arg Ser Tyr Arg Leu Ser Met Lys Val Lys Ala Phe	
	325	330 335
15	Pro Ser Pro Glu Val Val Trp Leu Lys Asp Gly Leu Pro Ala Thr Glu	
	340	345 350
	Lys Ser Ala Arg Tyr Leu Thr Arg Gly Tyr Ser Leu Ile Ile Lys Asp	
	355	360 365
20	Val Thr Glu Glu Asp Ala Gly Asn Tyr Thr Ile Leu Leu Ser Ile Lys	
	370	375 380
	Gln Ser Asn Val Phe Lys Asn Leu Thr Ala Thr Leu Ile Val Asn Val	
25	385	390 395 400
	Lys Pro Gln Ile Tyr Glu Lys Ala Val Ser Ser Phe Pro Asp Pro Ala	
	405	410 415
30	Leu Tyr Pro Leu Gly Ser Arg Gln Ile Leu Thr Cys Thr Ala Tyr Gly	
	420	425 430
	Ile Pro Gln Pro Thr Ile Lys Trp Phe Trp His Pro Cys Asn His Asn	
	435	440 445

- 58 -

	His Ser Glu Ala Arg Cys Asp Phe Cys Ser Asn Asn Glu Glu Ser Phe
	450 455 460
5	Ile Leu Asp Ala Asp Ser Asn Met Gly Asn Arg Ile Glu Ser Ile Thr
	465 470 475 480
	Gln Arg Met Ala Ile Ile Glu Gly Lys Asn Lys Met Ala Ser Thr Leu
10	485 490 495
	Val Val Ala Asp Ser Arg Ile Ser Gly Ile Tyr Ile Cys Ile Ala Ser
	500 505 510
15	Asn Lys Val Gly Thr Val Gly Arg Asn Ile Ser Phe Tyr Ile Thr Asp
	515 520 525
	Val Pro Asn Gly Phe His Val Asn Leu Glu Lys Met Pro Thr Glu Gly
	530 535 540
20	Glu Asp Leu Lys Leu Ser Cys Thr Val Asn Lys Phe Leu Tyr Arg Asp
	545 550 555 560
	Val Thr Trp Ile Leu Leu Arg Thr Val Asn Asn Arg Thr Met His Tyr
25	565 570 575
	Ser Ile Ser Lys Gln Lys Met Ala Ile Thr Lys Glu His Ser Ile Thr
	580 585 590
	Leu Asn Leu Thr Ile Met Asn Val Ser Leu Gln Asp Ser Gly Thr Tyr
30	595 600 605
	Ala Cys Arg Ala Arg Asn Val Tyr Thr Gly Glu Glu Ile Leu Gln Lys
	610 615 620

- 59 -

Lys Glu Ile Thr Ile Arg Gly Glu His Cys Asn Lys Lys Ala Val Phe
625 630 635 640

5

Ser Arg Ile Ser Lys Phe Lys Ser Thr Arg Asn Asp Cys Thr Thr Gln
645 650 655

Ser Asn Val Lys His
660

10

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 668 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

25

Ser Glu Gln Asn Met Gln Ser Lys Val Leu Leu Ala Val Ala Leu Trp
1 5 10 15

Leu Cys Val Glu Thr Arg Ala Ala Ser Val Gly Leu Pro Ser Val Ser
20 25 30

30

Leu Asp Leu Pro Arg Leu Ser Ile Gln Lys Asp Ile Leu Thr Ile Lys
35 40 45

- 60 -

Ala Asn Thr Thr Leu Gln Ile Thr Cys Arg Gly Gln Arg Asp Leu Asp
 50 55 60

5 Trp Leu Trp Pro Asn Asn Gln Ser Gly Ser Glu Gln Arg Val Glu Val
 65 70 75 80

Thr Glu Cys Ser Asp Gly Leu Phe Cys Lys Thr Leu Thr Ile Pro Lys
 85 90 95

10 Val Ile Gly Asn Asp Thr Gly Ala Tyr Lys Cys Phe Tyr Arg Glu Thr
 100 105 110

Asp Leu Ala Ser Val Ile Tyr Val Tyr Val Gln Asp Tyr Arg Ser Pro
 115 120 125

Phe Ile Ala Ser Val Ser Asp Gln His Gly Val Val Tyr Ile Thr Glu
 130 135 140

20 Asn Lys Asn Lys Thr Val Val Ile Pro Cys Leu Gly Ser Ile Ser Asn
 145 150 155 160

Leu Asn Val Ser Leu Cys Ala Arg Tyr Pro Glu Lys Arg Phe Val Pro
 165 170 175

25 Asp Gly Asn Arg Ile Ser Trp Asp Ser Lys Lys Gly Phe Thr Ile Pro
 180 185 190

Ser Tyr Met Ile Ser Tyr Ala Gly Met Val Phe Cys Glu Ala Lys Ile
 195 200 205

30 Asn Asp Glu Ser Tyr Gln Ser Ile Met Tyr Ile Val Val Val Val Gly
 210 215 220

- 61 -

	Tyr Arg Ile Tyr Asp Val Val Leu Ser Pro Ser His Gly Ile Glu Leu	
	225	230 235 240
5	Ser Val Gly Glu Lys Leu Val Leu Asn Cys Thr Ala Arg Thr Glu Leu	
	245	250 255
	Asn Val Gly Ile Asp Phe Asn Trp Glu Tyr Pro Ser Ser Lys His Gln	
	260	265 270
10	His Lys Lys Leu Val Asn Arg Asp Leu Lys Thr Gln Ser Gly Ser Glu	
	275	280 285
	Met Lys Lys Phe Leu Ser Thr Leu Thr Ile Asp Gly Val Thr Arg Ser	
15	290	295 300
	Asp Gln Gly Leu Tyr Thr Cys Ala Ala Ser Ser Gly Leu Met Thr Lys	
	305	310 315 320
20	Lys Asn Ser Thr Phe Val Arg Val His Glu Lys Pro Phe Val Ala Phe	
	325	330 335
	Gly Ser Gly Met Glu Ser Leu Val Glu Ala Thr Val Gly Glu Arg Val	
	340	345 350
25	Arg Ile Pro Ala Lys Tyr Leu Gly Tyr Pro Pro Pro Glu Ile Lys Trp	
	355	360 365
	Tyr Lys Asn Gly Ile Pro Leu Glu Ser Asn His Thr Ile Lys Ala Gly	
30	370	375 380
	His Val Leu Thr Ile Met Glu Val Ser Glu Arg Asp Thr Gly Asn Tyr	
	385	390 395 400

- 62 -

	Thr Val Ile Leu Thr Asn Pro Ile Ser Lys Glu Lys Gln Ser His Val	
	405	415
5	Val Ser Leu Val Val Tyr Val Pro Pro Gln Ile Gly Glu Lys Ser Leu	
	420	430
	Ile Ser Pro Val Asp Ser Tyr Gln Tyr Gly Thr Thr Gln Thr Leu Thr	
10	435	445
	Cys Thr Val Tyr Ala Ile Pro Pro Pro His His Ile His Trp Tyr Trp	
	450	460
15	Gln Leu Glu Glu Glu Cys Ala Asn Glu Pro Ser Gln Ala Val Ser Val	
	465	480
	Thr Asn Pro Tyr Pro Cys Glu Glu Trp Arg Ser Val Glu Asp Phe Gln	
	485	495
20	Gly Gly Asn Lys Ile Ala Val Asn Lys Asn Gln Phe Ala Leu Ile Glu	
	500	510
	Gly Lys Asn Lys Thr Val Ser Thr Leu Val Ile Gln Ala Ala Asn Val	
25	515	525
	Ser Ala Leu Tyr Lys Cys Glu Ala Val Asn Lys Val Gly Arg Gly Glu	
	530	540
30	Arg Val Ile Ser Phe His Val Thr Arg Gly Pro Glu Ile Thr Leu Gln	
	545	560
	Pro Asp Met Gln Pro Thr Glu Gln Glu Ser Val Ser Leu Trp Cys Thr	
	565	575

- 63 -

Ala Asp Arg Ser Thr Phe Glu Asn Leu Thr Trp Tyr Lys Leu Gly Pro
580 585 590

5 Gln Pro Leu Pro Ile His Val Gly Glu Leu Pro Thr Pro Val Cys Lys
595 600 605

Asn Leu Asp Thr Leu Trp Lys Leu Asn Ala Thr Met Phe Ser Asn Ser
610 615 620

10 Thr Asn Asp Ile Leu Ile Met Glu Leu Lys Asn Ala Ser Leu Gln Asp
625 630 635 640

Gln Gly Asp Tyr Val Cys Leu Ala Gln Asp Arg Lys Thr Lys Lys Arg
15 645 650 655

His Cys Val Val Arg Gln Leu Thr Val Leu Glu Arg
660 665

20 (2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 780 amino acids
(B) TYPE: amino acid
25 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

30

- 64 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

5	Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu Cys Ala Leu Leu Ser
	1 5 10 15
	Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly Ser Lys Leu Lys Asp Pro
	20 25 30
10	Glu Leu Ser Leu Lys Gly Thr Gln His Ile Met Gln Ala Gly Gln Thr
	35 40 45
	Leu His Leu Gln Cys Arg Gly Glu Ala Ala His Lys Trp Ser Leu Pro
15	50 55 60
	Glu Met Val Ser Lys Glu Ser Glu Arg Leu Ser Ile Thr Lys Ser Ala
	65 70 75 80
20	Cys Gly Arg Asn Gly Lys Gln Phe Cys Ser Thr Leu Thr Leu Asn Thr
	85 90 95
	Ala Gln Ala Asn His Thr Gly Phe Tyr Ser Cys Lys Tyr Leu Ala Val
	100 105 110
25	Pro Thr Ser Lys Lys Lys Glu Thr Glu Ser Ala Ile Tyr Ile Phe Ile
	115 120 125
	Ser Asp Thr Gly Arg Pro Phe Val Glu Met Tyr Ser Glu Ile Pro Glu
30	130 135 140
	Ile Ile His Met Thr Glu Gly Arg Glu Leu Val Ile Pro Cys Arg Val
	145 150 155 160

- 65 -

	Thr Ser Pro Asn Ile Thr Val Thr Leu Lys Lys Phe Pro Leu Asp Thr		
	165	170	175
5	Leu Ile Pro Asp Gly Lys Arg Ile Ile Trp Asp Ser Arg Lys Gly Phe		
	180	185	190
	Ile Ile Ser Asn Ala Thr Tyr Lys Glu Ile Gly Leu Leu Thr Cys Glu		
10	195	200	205
	Ala Thr Val Asn Gly His Leu Tyr Lys Thr Asn Tyr Leu Thr His Arg		
	210	215	220
15	Gln Thr Asn Thr Ile Ile Asp Val Gln Ile Ser Thr Pro Arg Pro Val		
	225	230	235 240
	Lys Leu Leu Arg Gly His Thr Leu Val Leu Asn Cys Thr Ala Thr Thr		
	245	250	255
20	Pro Leu Asn Thr Arg Val Gln Met Thr Trp Ser Tyr Pro Asp Glu Lys		
	260	265	270
	Asn Lys Arg Ala Ser Val Arg Arg Arg Ile Asp Gln Ser Asn Ser His		
25	275	280	285
	Ala Asn Ile Phe Tyr Ser Val Leu Thr Ile Asp Lys Met Gln Asn Lys		
	290	295	300
30	Asp Lys Gly Leu Tyr Thr Cys Arg Val Arg Ser Gly Pro Ser Phe Lys		
	305	310	315 320
	Ser Val Asn Thr Ser Val His Ile Tyr Asp Lys Ala Phe Ile Thr Val		
	325	330	335

- 66 -

Lys His Arg Lys Gln Gln Val Leu Glu Thr Val Ala Gly Lys Arg Ser
 340 345 350

5

Tyr Arg Leu Ser Met Lys Val Lys Ala Phe Pro Ser Pro Glu Val Val
 355 360 365

Trp Leu Lys Asp Gly Leu Pro Ala Thr Glu Lys Ser Ala Arg Tyr Leu
 370 375 380

10

Thr Arg Gly Tyr Ser Leu Ile Ile Lys Asp Val Thr Glu Glu Asp Ala
 385 390 395 400

Gly Asn Tyr Thr Ile Leu Leu Ser Ile Lys Gln Ser Asn Val Phe Lys
 405 410 415

15

Asn Leu Thr Ala Thr Leu Ile Val Asn Val Lys Pro Gln Ile Tyr Glu
 420 425 430

20

Lys Ala Val Ser Ser Phe Pro Asp Pro Ala Leu Tyr Pro Leu Gly Ser
 435 440 445

Arg Gln Ile Leu Thr Cys Thr Ala Tyr Gly Ile Pro Gln Pro Thr Ile
 450 455 460

25

Lys Trp Phe Trp His Pro Cys Asn His Asn His Ser Glu Ala Arg Cys
 465 470 475 480

Asp Phe Cys Ser Asn Asn Glu Glu Ser Phe Ile Leu Asp Ala Asp Ser
 485 490 495

30

Asn Met Gly Asn Arg Ile Glu Ser Ile Thr Gln Arg Met Ala Ile Ile
 500 505 510

- 67 -

	Glu Gly Lys Asn Lys Met Ala Ser Thr Leu Val Val Ala Asp Ser Arg
	515 520 525
5	Ile Ser Gly Ile Tyr Ile Cys Ile Ala Ser Asn Lys Val Gly Thr Val
	530 535 540
	Gly Arg Asn Ile Ser Phe Tyr Ile Thr Asp Val Pro Asn Gly Phe His
10	545 550 555 560
	Val Asn Leu Glu Lys Met Pro Thr Glu Gly Glu Asp Leu Lys Leu Ser
	565 570 575
15	Cys Thr Val Asn Lys Phe Leu Tyr Arg Asp Val Thr Trp Ile Leu Leu
	580 585 590
	Arg Thr Val Asn Asn Arg Thr Met His Tyr Ser Ile Ser Lys Gln Lys
	595 600 605
20	Met Ala Ile Thr Lys Glu His Ser Ile Thr Leu Asn Leu Thr Ile Met
	610 615 620
	Asn Val Ser Leu Gln Asp Ser Gly Thr Tyr Ala Cys Arg Ala Arg Asn
25	625 630 635 640
	Val Tyr Thr Gly Glu Glu Ile Leu Gln Lys Lys Glu Ile Thr Ile Arg
	645 650 655
30	Asp Gln Glu Ala Pro Tyr Leu Leu Arg Asn Leu Ser Asp His Thr Val
	660 665 670
	Ala Ile Ser Ser Ser Thr Thr Leu Asp Cys His Ala Asn Gly Val Pro
	675 680 685

- 68 -

Glu Pro Gln Ile Thr Trp Phe Lys Asn Asn His Lys Ile Gln Gln Glu
690 695 700

5 Pro Gly Ile Ile Leu Gly Pro Gly Ser Ser Thr Leu Phe Ile Glu Arg
705 710 715 720

Val Thr Glu Glu Asp Glu Gly Val Tyr His Cys Lys Ala Thr Asn Gln
725 730 735

10 Lys Gly Ser Val Glu Ser Ser Ala Tyr Leu Thr Val Gln Gly Thr Ser
740 745 750

15 Asp Lys Ser Asn Leu Glu Leu Ile Thr Leu Thr Cys Thr Cys Val Ala
755 760 765

Ala Thr Leu Phe Trp Leu Leu Leu Thr Leu Leu Ile
770 775 780

20 (2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 788 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

30

- 69 -

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:15:

5	Met Gln Ser Lys Val Leu Leu Ala Val Ala Leu Trp Leu Cys Val Glu
	1 5 10 15
	Thr Arg Ala Ala Ser Val Gly Leu Pro Ser Val Ser Leu Asp Leu Pro
	20 25 30
10	Arg Leu Ser Ile Gln Lys Asp Ile Leu Thr Ile Lys Ala Asn Thr Thr
	35 40 45
	Leu Gln Ile Thr Cys Arg Gly Gln Arg Asp Leu Asp Trp Leu Trp Pro
15	50 55 60
	Asn Asn Gln Ser Gly Ser Glu Gln Arg Val Glu Val Thr Glu Cys Ser
	65 70 75 80
20	Asp Gly Leu Phe Cys Lys Thr Leu Thr Ile Pro Lys Val Ile Gly Asn
	85 90 95
	Asp Thr Gly Ala Tyr Lys Cys Phe Tyr Arg Glu Thr Asp Leu Ala Ser
	100 105 110
25	Val Ile Tyr Val Tyr Val Gln Asp Tyr Arg Ser Pro Phe Ile Ala Ser
	115 120 125
	Val Ser Asp Gln His Gly Val Val Tyr Ile Thr Glu Asn Lys Asn Lys
30	130 135 140
	Thr Val Val Ile Pro Cys Leu Gly Ser Ile Ser Asn Leu Asn Val Ser
	145 150 155 160

- 70 -

	Leu Cys Ala Arg Tyr Pro Glu Lys Arg Phe Val Pro Asp Gly Asn Arg
	165 170 175
5	Ile Ser Trp Asp Ser Lys Lys Gly Phe Thr Ile Pro Ser Tyr Met Ile
	180 185 190
	Ser Tyr Ala Gly Met Val Phe Cys Glu Ala Lys Ile Asn Asp Glu Ser
	195 200 205
10	Tyr Gln Ser Ile Met Tyr Ile Val Val Val Val Gly Tyr Arg Ile Tyr
	210 215 220
	Asp Val Val Leu Ser Pro Ser His Gly Ile Glu Leu Ser Val Gly Glu
15	225 230 235 240
	Lys Leu Val Leu Asn Cys Thr Ala Arg Thr Glu Leu Asn Val Gly Ile
	245 250 255
20	Asp Phe Asn Trp Glu Tyr Pro Ser Ser Lys His Gln His Lys Lys Leu
	260 265 270
	Val Asn Arg Asp Leu Lys Thr Gln Ser Gly Ser Glu Met Lys Lys Phe
	275 280 285
25	Leu Ser Thr Leu Thr Ile Asp Gly Val Thr Arg Ser Asp Gln Gly Leu
	290 295 300
	Tyr Thr Cys Ala Ala Ser Ser Gly Leu Met Thr Lys Lys Asn Ser Thr
30	305 310 315 320
	Phe Val Arg Val His Glu Lys Pro Phe Val Ala Phe Gly Ser Gly Met
	325 330 335

- 71 -

	Glu Ser Leu Val Glu Ala Thr Val Gly Glu Arg Val Arg Ile Pro Ala	
	340	345 350
5	Lys Tyr Leu Gly Tyr Pro Pro Pro Glu Ile Lys Trp Tyr Lys Asn Gly	
	355	360 365
	Ile Pro Leu Glu Ser Asn His Thr Ile Lys Ala Gly His Val Leu Thr	
10	370	375 380
	Ile Met Glu Val Ser Glu Arg Asp Thr Gly Asn Tyr Thr Val Ile Leu	
	385	390 395 400
15	Thr Asn Pro Ile Ser Lys Glu Lys Gln Ser His Val Val Ser Leu Val	
	405	410 415
	Val Tyr Val Pro Pro Gln Ile Gly Glu Lys Ser Leu Ile Ser Pro Val	
	420	425 430
20	Asp Ser Tyr Gln Tyr Gly Thr Thr Gln Thr Leu Thr Cys Thr Val Tyr	
	435	440 445
	Ala Ile Pro Pro Pro His His Ile His Trp Tyr Trp Gln Leu Glu Glu	
25	450	455 460
	Glu Cys Ala Asn Glu Pro Ser Gln Ala Val Ser Val Thr Asn Pro Tyr	
	465	470 475 480
30	Pro Cys Glu Glu Trp Arg Ser Val Glu Asp Phe Gln Gly Gly Asn Lys	
	485	490 495
	Ile Ala Val Asn Lys Asn Gln Phe Ala Leu Ile Glu Gly Lys Asn Lys	
	500	505 510

- 72 -

Thr Val Ser Thr Leu Val Ile Gln Ala Ala Asn Val Ser Ala Leu Tyr
515 520 525

5 Lys Cys Glu Ala Val Asn Lys Val Gly Arg Gly Glu Arg Val Ile Ser
530 535 540

Phe His Val Thr Arg Gly Pro Glu Ile Thr Leu Gln Pro Asp Met Gln
545 550 555 560

10 Pro Thr Glu Gln Glu Ser Val Ser Leu Trp Cys Thr Ala Asp Arg Ser
565 570 575

Thr Phe Glu Asn Leu Thr Trp Tyr Lys Leu Gly Pro Gln Pro Leu Pro
15 580 585 590

Ile His Val Gly Glu Leu Pro Thr Pro Val Cys Lys Asn Leu Asp Thr
595 600 605

20 Leu Trp Lys Leu Asn Ala Thr Met Phe Ser Asn Ser Thr Asn Asp Ile
610 615 620

Leu Ile Met Glu Leu Lys Asn Ala Ser Leu Gln Asp Gln Gly Asp Tyr
625 630 635 640

25 Val Cys Leu Ala Gln Asp Arg Lys Thr Lys Lys Arg His Cys Val Val
645 650 655

Arg Gln Leu Thr Val Leu Glu Arg Val Ala Pro Thr Ile Thr Gly Asn
30 660 665 670

Leu Glu Asn Gln Thr Thr Ser Ile Gly Glu Ser Ile Glu Val Ser Cys
675 680 685

- 73 -

	Thr Ala Ser Gly Asn Pro Pro Pro Gln Ile Met Trp Phe Lys Asp Asn
	690 695 700
5	Glu Thr Leu Val Glu Asp Ser Gly Ile Val Leu Lys Asp Gly Asn Arg
	705 710 715 720
	Asn Leu Thr Ile Arg Arg Val Arg Lys Glu Asp Glu Gly Leu Tyr Cys
	725 730 735
10	Gln Ala Cys Ser Val Leu Gly Cys Ala Lys Val Glu Ala Phe Phe Ile
	740 745 750
	Ile Glu Gly Ala Gln Glu Lys Thr Asn Leu Glu Ile Ile Ile Leu Val
15	755 760 765
	Gly Thr Thr Val Ile Ala Met Phe Phe Trp Leu Leu Leu Val Ile Ile
	770 775 780
20	Leu Gly Thr Val
	785

(2) INFORMATION FOR SEQ ID NO:16:

- 25 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2264 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

- 74 -

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:16:

5	GGTGTGGTCG CTGCGTTTCC TCTGCCTGCG CCGGGCATCA CTTGCGCGCC GCAGAAAGTC	60
	CGTCTGGCAG CCTGGATATC CTCTCCTACC GGCACCCGCA GACGCCCTG CAGCCGCGGT	120
	CGGCGCCCGG GCTCCCTAGC CCTGTGCGCT CAACTGTCCT GCGCTGCGGG GTGCCGCGAG	180
10	TTCCACCTCC GCGCCTCCTT CTCTAGACAG GCGCTGGGAG AAAGAACCGG CTCCCGAGTT	240
	CCGGCATTTC GCCCGGCTCG AGGTGCAGGA TGCAGAGCAA GGTGCTGCTG GCCGTCGCCC	300
15	TGTGGCTCTG CGTGGAGACC CGGGCCGCCT CTGTGGGTTT GCCTAGTGTT TCTCTTGATC	360
	TGCCCAGGCT CAGCATACAA AAAGACATAC TTACAATTAA GGCTAATACA ACTCTTCAAA	420
	TTACTTGCAg GGGACAGAGG GACTTGGACT GGCTTTGGCC CAATAATCAG AGTGGCAGTG	480
20	AGCAAAGGGT GGAGGTGACT GAGTGCAGCG ATGGCCTCTT CTGTAAGACA CTCACAATTC	540
	CAAAAGTGAT CGGAAATGAC ACTGGAGCCT ACAAGTGCTT CTACCGGGAA ACTGACTTGG	600
25	CCTCGGTCAT TTATGTCTAT GTTCAAGATT ACAGATCTCC ATTTATTGCT TCTGTTAGTG	660
	ACCAACATGG AGTCGTGTAC ATTACTGAGA ACAAAAACAA AACTGTGGTG ATTCCATGTC	720
	TCGGGTCCAT TTCAAATCTC AACGTGTAC TTTGTGCAAG ATACCCAGAA AAGAGATTG	780
30	TTCTGATGG TAACAGAATT TCCTGGGACA GCAAGAAGGG CTTTACTATT CCCAGCTACA	840
	TGATCAGCTA TGCTGGCATG GTCTTCTGTG AAGCAAAAAT TAATGATGAA AGTTACCACT	900

- 75 -

CTATTATGTA CATAGTTGTC GTTGTAGGGT ATAGGATTTA TGATGTGGTT CTGAGTCCGT 960

5 CTATGGAAT TGAATATCT GTTGGAGAAA AGCTTGCTT AAATTGTACA GCAAGAACTG 1020

AACTAAATGT GGGGATTGAC TTCAACTGGG AATACCCTTC TTCGAAGCAT CAGCATAAGA 1080

AACTGTAAA CCGAGACCTA AAAACCCAGT CTGGGAGTGA GATGAAGAAA TTTTGTAGCA 1140

10 CCTTAATAT AGATGGTGA ACCCGGAGTG ACCAAGGATT GTACACCTGT GCAGCATCCA 1200

GTGGGCTGAT GACCAAGAAG AACAGCACAT TTGTCAGGGT CCATGAAAAA CCTTTTGTG 1260

15 CTTTGTGAAG TGGCATGGAA TCTCTGGTG AAGCCACGGT GGGGAGCGT GTCAGAATCC 1320

CTGCGAAGTA CCTTGTTAC CCACCCAG AAATAAATG GTATAAAT GGAATACCC 1380

TTGAGTCAA TCACACAATT AAAGCGGGC ATGTACTGAC GATTATGGAA GTGAGTAAA 1440

20 GAGACACAGG AAATTACACT GTCATCTTA CCAATCCAT TTCAAAGGAG AAGCAGAGCC 1500

ATGTGGTCTC TCTGGTTGTG TATGTCCAC CCCAGATTG TGAGAAATCT CTAATCTCTC 1560

25 CTGTGGATTC CTACCAGTAC GGCACCACTC AAACGCTGAC ATGTACGGTC TATGCCATTC 1620

CTCCCCGCA TCACATCCAC TGGTATTGGC AGTTGGAGGA AGAGTGGCC AACGAGCCCA 1680

GCCAAGCTGT CTCAGTGACA AACCCATACC CTTGTGAAGA ATGGAGAAST GTGGAGGACT 1740

30 TCCAGGGAGG AAATAAAT GCGTTAATA AAAATCAATT TGCTCTAATT GAAGGAAAAA 1800

ACAAAATGT AAGTACCCTT GTTATCCAAG CGGCAATGT GTCAGCTTTG TACAAATGTG 1860

AAGCGGTCAA CAAAGTCGGG AGAGGAGAGA GGGTGATCTC CTTCCACGTG ACCAGGGGTC 1920

- 76 -

5 CTGAAATTAC TTTGCAACCT GACATGCAGC CCACTGAGCA GGAGAGCGTG TCTTTGTGGT 1980
GCACTGCAGA CAGATCTACG TTTGAGAACC TCACATGGTA CAAGCTTGGC CCACAGCCTC 2040
TGCCAATCCA TGTGGGAGAG TTGCCCACAC CTGTTTGCAA GAACTTGGAT ACTCTTTGGA 2100
AATTGAATGC CACCATGTTT TCTAATAGCA CAAATGACAT TTTGATCATG GAGCTTAAGA 2160
10 ATGCATCCTT GCAGGACCAA GGAGACTATG TCTGCCTTGC TCAAGACAGG AAGACCAAGA 2220
AAAGACATTG CGTGGTCAGG CAGCTCACAG TCCTAGAGCG TTAA 2264

(2) INFORMATION FOR SEQ ID NO:17:

15

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2352 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

20

(D) TOPOLOGY: linear

(11) MOLECULE TYPE: DNA (genomic)

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

60 GCGTCACCA TGGTCAGTA CTGGGACACC GGGGTCCTGC TGTGCGCGCT GCTCAGCTGT
30 CTGCTTCTCA CAGGATCTAG TTCAGGTTCA AAATTAAGG ATCCTGAACT GAGTTTAAAA 120
GGCACCCAGC ACATCATGCA AGCAGGCCAG ACACTGCATC TCCAATGCAG GGGGGAAGCA 180
GCCATAAAT GGTCTTTGCC TGAAATGCTG AGTAAGGAAA GCGAAAGGCT GAGCATAACT 240

- 77 -

	AAATCTGCCT GTGGAAGAAA TGGCAAACAA TTCTGCAGTA CTTTAACTT GAACACAGCT	300
5	CAAGCAAACC ACACTGGCTT CTACAGCTGC AAATATCTAG CTGTACCTAC TTCAAAGAAG	360
	AAGGAAACAG AATCTGCAAT CTATATATTT ATTAGTGATA CAGGTAGACC TTTCGTAGAG	420
	ATGTACAGTG AAATCCCCGA AATTATACAC ATGACTGAAG GAAGGGAGCT CGTCATTCCC	480
10	TGCCGGGTTA CGTCACCTAA CATCACTGTT ACTTTAAAAA AGTTTCCACT TGACACTTTG	540
	ATCCCTGATG GAAAACGCAT AATCTGGGAC AGTAGAAAGG GCTTCATCAT ATCAAATGCA	600
15	ACGTACAAAG AAATAGGGCT TCTGACCTGT GAAGCAACAG TCAATGGGCA TTTGTATAAG	660
	ACAACTATC TCACACATCG ACAAACCAAT ACAATCATAG ATGTCCAAAT AAGCACACCA	720
	CGCCCACTCA AATTACTTAG AGGCCATACT CTTGTCTCA ATTGTACTGC TACCACTCCC	780
20	TTGAACACGA GAGTTCAAAT GACCTGGAGT TACCCTGATG AAAAAATAA GAGAGCTTCC	840
	GTAAGGCGAC GAATTGACCA AAGCAATTCC CATGCCAACA TATTCTACAG TGTCTTACT	900
25	ATTGACAAAA TGCAGAACAA AGACAAAGGA CTTTATACTT GTCGTGTAAG GAGTGGACCA	960
	TCATTCAAAT CTGTTAACAC CTCAGTGCAT ATATATGATA AAGCATTCACT CACTGTGAAA	1020
	CATCGAAAAC AGCAGGTGCT TGAACCCTA GCTGGCAAGC GGTCTTACCG GCTCTCTATG	1080
30	AAAGTGAAGG CATTTCCTTC GCCGGAAGTT STATGGTTAA AAGATGGGTT ACCTGCGACT	1140
	GAGAAATCTG CTCGCTATTT GACTCGTGGC TACTCGTTAA TTATCAAGGA CGTAACTGAA	1200
	GAGGATSCAG GGAATTATAC AATCTTGCTG AGCATAAAAC AGTCAAATGT GTTTAAAAAC	1260

- 78 -

	CTCACTGCCA CTCTAATTGT CAATGTGAAA CCCAGATT ACGAAAAGGC CGTGTCTCG	1320
5	TTCCAGACC CGGCTCTCTA CCCACTGGG AGCAGACAAA TCCTGACTTG TACCGCATAT	1380
	GGTATCCCTC AACCTACAAT CAAGTGGTC TGGCACCCCT GTAACCATAA TCATTCCGAA	1440
	GCAAGGTGTG ACTTTTGTTT CAATAATGAA GAGTCCTTGA TCCTGGATGC TGACAGCAAC	1500
10	ATGGGAAACA GAATTGAGAG CATCACTCAG CGCATGGCAA TAATAGAAGG AAAGAATAAG	1560
	ATGGCTAGCA CCTTGGTTGT GGCTGACTCT AGAATTTCTG GAATCTACAT TTGCATAGCT	1620
15	TCCAATAAG TTGGGACTGT GGAAGAAAC ATAAGCTTTT ATATCACAGA TGTGCCAAAT	1680
	GGGTTTCATG TTAACCTGGA AAAATGCCG ACGGAAGGAG AGGACCTGAA ACTGTCTTGC	1740
	ACAGTTAACA AGTTCTTATA CAGAGAGTTT ACTTGGATTT TACTGCGGAC AGTTAATAAC	1800
20	AGAACAATGC ACTACAGTAT TAGCAAGCAA AAAATGGCCA TCACTAAGGA GCACTCCATC	1860
	ACTCTTAATC TTACCATCAT GAATGTTTCC CTGCAAGATT CAGGCACCTA TGCCTGCAGA	1920
25	GCCAGGAATG TATACACAGG GGAAGAAATC CTCCAGAAGA AAGAAATTAC AATCAGAGAT	1980
	CAGGAAGCAC CATACCTCCT GCGAAACCTC AGTGATCACA CAGTGGCCAT CAGCAGTTCC	2040
	ACCACTTTAG ACTGTCTATG TAATGGTGTG CCCGAGCCTC AGATCACTTG GTTTAAAAAC	2100
30	AACCACAAAA TACAACAAGA GCCTGGAATT ATTTTAGGAC CAGGAAGCAG CACGCTGTTT	2160
	ATTGAAAGAG TCACAGAAGA GGATGAAGGT GTCTATCACT GCAAAGCCAC CAACCAGAAG	2220
	GGCTCTGTGG AAAGTTCAGC ATACCTCACT GTTCAAGGAA CCTCGGACAA GTCTAATCTG	2280

- 79 -

GAGCTGATCA CTCTAACATG CACCTGTGTG GCTGCGACTC TCTTCTGGCT CCTATTAACC 2340

CTCCTTATCT AA 2352

5

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 2383 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

20

CTCGAGGTGC AGGATGCAGA GCAAGGTGCT GCTGGCCGTC GCCCTGTGGC TCTGCGTGGG 60

GACCCGGGCC GCCTCTGTGG GTTTCCTAG TGTTCCTTT GATCTGCCA GGCTCAGCAT 120

25

ACAAAAAGAC ATACTTACAA TTAAGGCTAA TACAACCTTT CAAATTACTT GCAGGGGACA 180

GAGGGACTTG GACTGGCTTT GGCCCAATAA TCAGAGTGGC AGTGAGCAAA GGGTGGAGGT 240

GACTGAGTGC AGCGATGGCC TCTTCTGTAA GACACTCACA ATTCCAAAAG TGATCGGAAA 300

30

TGACACTGGA GCCTACAAGT GCTTCTACCG GGAAACTGAC TTGGCCTCGG TCATTATGT 360

CTATGTTCAA GATTACAGAT CTCCATTAT TGCTTCTGTT AGTGACCAAC ATGGAGTCGT 420

GTACATTACT GAGAACAAAA ACAAACTGT GGTGATTCCA TGTCTCGGT CCATTTCAAA 480

- 80 -

5
10
15
20
25
30

TCTCAACGTG TCACCTTTGTG CAAGATACCC AGAAAAGAGA TTTGTCTCTG ATGGTAACAG 540
AATTCCTGG GACAGCAAGA AGGGCTTTAC TATCCCAGC TACATGATCA GCTATGCTGG 600
CATGGTCTTC TGTGAAGCAA AAATTAATGA TGAAAGTTAC CAGTCTATTA TGTACATAGT 660
TGTCGTTGTA GGGTATAGGA TTTATGATGT GGTCTGAGT CCGTCTCATG GAATTGAACT 720
ATCTGTTGGA GAAAAGCTTG TCTTAAATTG TACAGCAAGA ACTGAACTAA ATGTGGGGAT 780
TGACTTCAAC TGGGAATACC CTTCTTCGAA GCATCAGCAT AAGAACTTG TAAACCGAGA 840
CCTAAAAACC CAGTCTGGGA GTGAGATGAA GAAATTTTGG AGCACCTTAA CTATAGATGG 900
TGTAACCCGG AGTGACCAAG GATTGTACAC CTGTGCAGCA TCCAGTGGGC TGATGACCAA 960
GAAGAACAGC ACATTGTCA GGGTCCATGA AAAACCTTTT GTTGCTTTTG GAAGTGGCAT 1020
GGAATCTCTG GTGGAAGCCA CGGTGGGGGA GCGTGTGAGA ATCCCTGCCA AGTACCTTGG 1080
TTACCCACCC CCAGAAATAA AATGGTATAA AAATGGAATA CCCCTTGAGT CCAATCACAC 1140
AATTAAAGCG GGGCATGTAC TGACGATTAT GGAAGTGAGT GAAAGAGACA CAGGAAATTA 1200
CACTGTCTC CTTACCAATC CCATTTCAA GGAGAAGCAG AGCCATGTGG TCTCTCTGGT 1260
TGTGTATGTC CCACCCGAGA TTGGTGAGAA ATCTCTAATC TCTCTGTGG ATCTCTACCA 1320
GTACGGCACC ACTCAAACGC TGACATGTAC GGTCTATGCC ATTCCTCCCC CGCATCACAT 1380
CCACTGGTAT TGGCASTTGG AGGAAGAGTG CGCCAACGAG CCCAGCCAAG CTGTCTCAGT 1440
GACAAACCCA TACCCTTGTG AAGAATGGAG AAGTGTGGAG GACTTCCAGG GAGGAAATAA 1500

- 81 -

	AATTGCCGTT AATAAAAAATC AATTTGCTCT AATTGAAGGA AAAAACAAAA CTGTAAGTAC	1560
5	CCTTGTTATC CAAGCGGCAA ATGTGTCAGC TTTGTACAAA TGTGAAGCGG TCAACAAAGT	1620
	CGGGAGAGGA GAGAGGGTGA TCTCCTTCCA CGTGACCAGG GGTCTGAAA TTACTTTGCA	1680
	ACCTGACATG CAGCCCACTG AGCAGGAGAG CGTGTCTTTG TGGTCACTG CAGACAGATC	1740
10	TACGTTTGAG AACCTCACAT GGTACAAGCT TGGCCACAG CCTCTGCCAA TCCATGTGGG	1800
	AGAGTTGCCC ACACCTGTTT GCAAGAACTT GGATACTCTT TGGAAATTGA ATGCCACCAT	1860
15	GTCTCTAAT AGCACAAATG ACATTTTGAT CATGGAGCTT AAGAATGCAT CCTGCAGGA	1920
	CCAAGGAGAC TATGTCTGCC TTGCTCAAGA CAGGAAGACC AAGAAAAGAC ATTGCGTGGT	1980
	CAGGCAGCTC ACAGTCCTAG AGCGTGTGGC ACCCAGGATC ACAGGAAACC TGGAGAATCA	2040
20	GACGACAAGT ATTGGGGAAA GCATCGAAGT CTCATGCACG GCATCTGGGA ATCCCCCTCC	2100
	ACAGATCATG TGGTTTAAAG ATAATGAGAC CCTGTAGAA GACTCAGGCA TTGTATTGAA	2160
25	GGATGGGAAC CGGAACCTCA CTATCCGAG AGTGAGGAAG GAGGACGAAG GCCTCTACAC	2220
	CTGCCAGGCA TGCAGTGTTT TTGGCTGTGC AAAAGTGGAG GCATTTTCA TAATAGAAGG	2280
	TGCCCAGGAA AAGACGAACT TGGAAATCAT TATTCTAGTA GGCACGACGG TGATTGCCAT	2340
30	GTCTCTCTGG CTACTTCTTG TCATCATCCT AGGGACCGTT TAA	2383

- 82 -

WHAT IS CLAIMED IS:

1. A soluble VEGF inhibitor in substantially pure form
5 which specifically binds VEGF and inhibits cellular VEGF
receptor activity.

2. The soluble VEGF inhibitor according to Claim 1
wherein the soluble VEGF receptor is selected from the
10 group consisting of sVEGF-RI, sVEGF-RII, sVEGF-RTMI and
sVEGF-RTMII.

3. The soluble VEGF inhibitor of Claim 2 corresponding
to sVEGF-RI comprising the amino acid sequence:
15

Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu

Cys Ala Leu Leu Ser Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly

20 Ser Lys Leu Lys Asp Pro Glu Leu Ser Leu Lys Gly Thr Gln His

Ile Met Gln Ala Gly Gln Thr Leu His Leu Gln Cys Arg Gly Glu

25 Ala Ala His Lys Trp Ser Leu Pro Glu Met Val Ser Lys Glu Ser

Glu Arg Leu Ser Ile Thr Lys Ser Ala Cys Gly Arg Asn Gly Lys

30

- 83 -

Gln Phe Cys Ser Thr Leu Thr Leu Asn Thr Ala Gln Ala Asn His
5 Thr Gly Phe Tyr Ser Cys Lys Tyr Leu Ala Val Pro Thr Ser Lys
Lys Lys Glu Thr Glu Ser Ala Ile Tyr Ile Phe Ile Ser Asp Thr
Gly Arg Pro Phe Val Glu Met Tyr Ser Glu Ile Pro Glu Ile Ile
10 His Met Thr Glu Gly Arg Glu Leu Val Ile Pro Cys Arg Val Thr
Ser Pro Asn Ile Thr Val Thr Leu Lys Lys Phe Pro Leu Asp Thr
Leu Ile Pro Asp Gly Lys Arg Ile Ile Trp Asp Ser Arg Lys Gly
15 Phe Ile Ile Ser Asn Ala Thr Tyr Lys Glu Ile Gly Leu Leu Thr
Cys Glu Ala Thr Val Asn Gly His Leu Tyr Lys Thr Asn Tyr Leu
20 Thr His Arg Gln Thr Asn Thr Ile Ile Asp Val Gln Ile Ser Thr

25

30

- 84 -

Pro Arg Pro Val Lys Leu Leu Arg Gly His Thr Leu Val Leu Asn
Cys Thr Ala Thr Thr Pro Leu Asn Thr Arg Val Gln Met Thr Trp
5 Ser Tyr Pro Asp Glu Lys Asn Lys Arg Ala Ser Val Arg Arg Arg
Ile Asp Gln Ser Asn Ser His Ala Asn Ile Phe Tyr Ser Val Leu
10 Thr Ile Asp Lys Met Gln Asn Lys Asp Lys Gly Leu Tyr Thr Cys
Arg Val Arg Ser Gly Pro Ser Phe Lys Ser Val Asn Thr Ser Val
His Ile Tyr Asp Lys Ala Phe Ile Thr Val Lys His Arg Lys Gln
15 Gln Val Leu Glu Thr Val Ala Gly Lys Arg Ser Tyr Arg Leu Ser
Met Lys Val Lys Ala Phe Pro Ser Pro Glu Val Val Trp Leu Lys
20 Asp Gly Leu Pro Ala Thr Glu Lys Ser Ala Arg Tyr Leu Thr Arg

25

30

- 85 -

Gly Tyr Ser Leu Ile Ile Lys Asp Val Thr Glu Glu Asp Ala Gly
Asn Tyr Thr Ile Leu Leu Ser Ile Lys Gln Ser Asn Val Phe Lys
5 Asn Leu Thr Ala Thr Leu Ile Val Asn Val Lys Pro Gln Ile Tyr
Glu Lys Ala Val Ser Ser Phe Pro Asp Pro Ala Leu Tyr Pro Leu
10 Gly Ser Arg Gln Ile Leu Thr Cys Thr Ala Tyr Gly Ile Pro Gln
Pro Thr Ile Lys Trp Phe Trp His Pro Cys Asn His Asn His Ser
Glu Ala Arg Cys Asp Phe Cys Ser Asn Asn Glu Glu Ser Phe Ile
15 Leu Asp Ala Asp Ser Asn Met Gly Asn Arg Ile Glu Ser Ile Thr
Gln Arg Met Ala Ile Ile Glu Gly Lys Asn Lys Met Ala Ser Thr
20 Leu Val Val Ala Asp Ser Arg Ile Ser Gly Ile Tyr Ile Cys Ile

25

30

- 86 -

Ala Ser Asn Lys Val Gly Thr Val Gly Arg Asn Ile Ser Phe Tyr
Ile Thr Asp Val Pro Asn Gly Phe His Val Asn Leu Glu Lys Met
5 Pro Thr Glu Gly Glu Asp Leu Lys Leu Ser Cys Thr Val Asn Lys
Phe Leu Tyr Arg Asp Val Thr Trp Ile Leu Leu Arg Thr Val Asn
10 Asn Arg Thr Met His Tyr Ser Ile Ser Lys Gln Lys Met Ala Ile
Thr Lys Glu His Ser Ile Thr Leu Asn Leu Thr Ile Met Asn Val
Ser Leu Gln Asp Ser Gly Thr Tyr Ala Cys Arg Ala Arg Asn Val
15 Tyr Thr Gly Glu Glu Ile Leu Gln Lys Lys Glu Ile Thr Ile Arg
Gly Glu His Cys Asn Lys Lys Ala Val Phe Ser Arg Ile Ser Lys
20 Phe Lys Ser Thr Arg Asn Asp Cys Thr Thr Gln Ser Asn Val Lys
His. (SEQ. ID. NO.: 6)

25

30

- 87 -

4. The soluble VEGF inhibitor of Claim 2 corresponding to sVEGF-RI comprising the amino acid sequence:

5 Ser Lys Leu Lys Asp Pro Glu Leu Ser Leu Lys Gly Thr Gln His

 Ile Met Gln Ala Gly Gln Thr Leu His Leu Gln Cys Arg Gly Glu

 Ala Ala His Lys Trp Ser Leu Pro Glu Met Val Ser Lys Glu Ser
10 Glu Arg Leu Ser Ile Thr Lys Ser Ala Cys Gly Arg Asn Gly Lys

 Gln Phe Cys Ser Thr Leu Thr Leu Asn Thr Ala Gln Ala Asn His

15 Thr Gly Phe Tyr Ser Cys Lys Tyr Leu Ala Val Pro Thr Ser Lys

 Lys Lys Glu Thr Glu Ser Ala Ile Tyr Ile Phe Ile Ser Asp Thr

 Gly Arg Pro Phe Val Glu Met Tyr Ser Glu Ile Pro Glu Ile Ile
20 His Met Thr Glu Gly Arg Glu Leu Val Ile Pro Cys Arg Val Thr

25

30

- 88 -

Ser Pro Asn Ile Thr Val Thr Leu Lys Lys Phe Pro Leu Asp Thr
Leu Ile Pro Asp Gly Lys Arg Ile Ile Trp Asp Ser Arg Lys Gly
5 Phe Ile Ile Ser Asn Ala Thr Tyr Lys Glu Ile Gly Leu Leu Thr
Cys Glu Ala Thr Val Asn Gly His Leu Tyr Lys Thr Asn Tyr Leu
10 Thr His Arg Gln Thr Asn Thr Ile Ile Asp Val Gln Ile Ser Thr
Pro Arg Pro Val Lys Leu Leu Arg Gly His Thr Leu Val Leu Asn
Cys Thr Ala Thr Thr Pro Leu Asn Thr Arg Val Gln Met Thr Trp
15 Ser Tyr Pro Asp Glu Lys Asn Lys Arg Ala Ser Val Arg Arg Arg
Ile Asp Gln Ser Asn Ser His Ala Asn Ile Phe Tyr Ser Val Leu
20 Thr Ile Asp Lys Met Gln Asn Lys Asp Lys Gly Leu Tyr Thr Cys

25

30

- 89 -

Arg Val Arg Ser Gly Pro Ser Phe Lys Ser Val Asn Thr Ser Val
His Ile Tyr Asp Lys Ala Phe Ile Thr Val Lys His Arg Lys Gln
5 Gln Val Leu Glu Thr Val Ala Gly Lys Arg Ser Tyr Arg Leu Ser
Met Lys Val Lys Ala Phe Pro Ser Pro Glu Val Val Trp Leu Lys
10 Asp Gly Leu Pro Ala Thr Glu Lys Ser Ala Arg Tyr Leu Thr Arg
Gly Tyr Ser Leu Ile Ile Lys Asp Val Thr Glu Glu Asp Ala Gly
Asn Tyr Thr Ile Leu Leu Ser Ile Lys Gln Ser Asn Val Phe Lys
15 Asn Leu Thr Ala Thr Leu Ile Val Asn Val Lys Pro Gln Ile Tyr
Glu Lys Ala Val Ser Ser Phe Pro Asp Pro Ala Leu Tyr Pro Leu
20 Gly Ser Arg Gln Ile Leu Thr Cys Thr Ala Tyr Gly Ile Pro Gln

25

30

- 90 -

Pro Thr Ile Lys Trp Phe Trp His Pro Cys Asn His Asn His Ser
Glu Ala Arg Cys Asp Phe Cys Ser Asn Asn Glu Glu Ser Phe Ile
5 Leu Asp Ala Asp Ser Asn Met Gly Asn Arg Ile Glu Ser Ile Thr
Gln Arg Met Ala Ile Ile Glu Gly Lys Asn Lys Met Ala Ser Thr
10 Leu Val Val Ala Asp Ser Arg Ile Ser Gly Ile Tyr Ile Cys Ile
Ala Ser Asn Lys Val Gly Thr Val Gly Arg Asn Ile Ser Phe Tyr
Ile Thr Asp Val Pro Asn Gly Phe His Val Asn Leu Glu Lys Met
15 Pro Thr Glu Gly Glu Asp Leu Lys Leu Ser Cys Thr Val Asn Lys
Phe Leu Tyr Arg Asp Val Thr Trp Ile Leu Leu Arg Thr Val Asn
20 Asn Arg Thr Met His Tyr Ser Ile Ser Lys Gln Lys Met Ala Ile

25

30

- 91 -

Thr Lys Glu His Ser Ile Thr Leu Asn Leu Thr Ile Met Asn Val

Ser Leu Gln Asp Ser Gly Thr Tyr Ala Cys Arg Ala Arg Asn Val

5

Tyr Thr Gly Glu Glu Ile Leu Gln Lys Lys Glu Ile Thr Ile Arg

Gly Glu His Cys Asn Lys Lys Ala Val Phe Ser Arg Ile Ser Lys

10

Phe Lys Ser Thr Arg Asn Asp Cys Thr Thr Gln Ser Asn Val Lys

His. (SEQ. ID. NO.: 12)

5. The soluble VEGF inhibitor of Claim 2 corresponding
15 to sVEGF-RII comprising the amino acid sequence:

MQSKVLLAVALWLCVETRAASVGLPSVSLDLPRLSIQKDILTIKANTTLQITCRGQR
DLDWLWPNNQSGSEQRVEVTECSDGLFCKTLTIPKVIGNDTGAYKCFYRETDLASVI
YVYVQDYRSPFIASVSDQHGCVYITENKNTVVIPCLGSISNLNVSLCARYPEKRFV
20 PDGNRISWDSKKGFTIPSYMISYAGMVFEAKINDESYQSIMYIVVVVGYRIYDVVL
SPSHGIELSVGEKLVNCTARTELNVGIDFNWEYPSSKHQHKLVNRDLKTQSGSEM
KKFLSTLTIDGVTRSDQGLYTCAASSGLMTKKNSTFVRVHEKPFVAFGSGMESLVEA
TVGERVRIPAKYLGYPPEIKWYKNGIPLESNHTIKAGHVLTIMEVSEKDTGNYTVI
LTNPISKEKQSHVSVLVVYVPPQIGEKSLISPVDSYQYGTQTTLCTVYAIPPPHEI
25 HWYWQLEEECANEPSQAVSVTNYPCEEWSVEDFQGGNKIAVNKNQFALIEGKNKT
VSTLVIQAANVSALYKCEAVNKVGRGERVISFEVTRGPEITLQPDMPTEQESVSLW
CTADRSTFENLTWYKLGPPQLPIHVGELEPTPVCKNLDLWKLNATMFSNSTNDILIM
ELKNASLQDQGDYVCLAQDRKTKKRHCVVRLTVLER. (SEQ.ID.NO.: 13)

30

- 92 -

6. The soluble VEGF inhibitor of Claim 2 corresponding to sVEGF-RTMI comprising the amino acid sequence:

5 MVSYWDTGVLLCALLSCLLLTGSSSSGSKLDPFELSLKGTQHIMQAGQTLHLQCRGEA
AHKWSLPPEMVSKESERLSITKSACGRNGKQFCSTLTLNTAQANHTGFYSCKYLAVPT
SKKETESAIYIFISDTGRPFVEMYSEIPEIIHMTGRELVIPCRVTSPNITVTLKK
FPLDTLIPDGKRIIWDSRKGFIISNATYKEIGLLTCEATVNGHLYKTNLTHRQTNT
IIDVQISTPRPVKLLRGHTLVLNCTATTPLNTRVQMTWSYPDEKNKRASVRRRIDQS
10 NSHANIFYSVLTIDKMQNKDKGLYTCRVRSGPSFKSVNTSVHIYDKAFITVKHRKQQ
VLETVAGKRSYRLSMKVKAFFPSPEVVWLKDGLPATEKSARYLTRGYSLLIKDVTEED
AGNYTILLSIKQSNVFNLTATLIVNVKPKIYEKAVSSFDPALYPLGSRQILTCTA
YGIPQPTIKWFHPCNHNHSEARCDFCSNNEESFILDADSNMGNRIESITQRMATIE
GKNKMASTLVVADSRISGIYICIASNKVGTVGRNISFYITDVPNGFHVNLKMPTEG
15 EDLKLSTVKNFLYRDVTWILLRTVNNRTMHYSISKQKMAITKEHSITLNLTIMNVS
LQDSGTIACRARNVYTGEIILQKKEITIRDQEAPYLLRNLSDHTVAISSSTTLTDLCHA
NGVPEPQITWFKNNHKIQEPGIILGPGSSSTLFIERVTEEDGVYHCKATNQKGSVE
SSAYLTVQGTSDKSNLELITLTCTCVAATLFWLLLTLLI. (SEQ. ID. NO.:
14)

20

7. The soluble VEGF inhibitor of Claim 2 corresponding to sVEGF-RTMII comprising the amino acid sequence:

MQSKVLLAVALWLCVETRAASVGLPSVSLDLPRLSIQKDILTILKANTTLQITCRGQR
25 DLDWLWPNNQSGSEQRVEVTECSDFLCKTLTIPKVIGNDTGAYKCFYRETDLASVI
YVYVQDYRSPFIASVSDQHGVVYITENKNKTVPCLGSISNLNVSLCARYPEKRFV
PDGNRISWDSKKGFTIPSYMISYAGMVCFEAKINDESYQSIMYIVVVVGYRIYDVVL
SPSHGIELSVGEKLVNLCTARTELNVGIDFNWEYPSKHKQHKLVNRDLKTQSGSEM
KKFLSTLTIDGVTRSDQGLYTCAASSGLMTKKNSTFVRVHEKPFVAFGSGMESLVEA
30 TVGERVRIPAKYLGYPPEIKWYKNGIPLESNETIKAGHVLTIMEVSRDTGNYTVI
LTNPISKEKQSHVVS LVVYVPPQIGEKSLI SPVDSYQYGTQTTLTCTVYAIPPPHHI

- 93 -

HWYWQLEEECANEPSQAVSVTNPYPCEEWRSVEDFQGGNKIAVNKNQFALIEGKNKT
VSTLVIQAANVSALYKCEAVNKVGRGERVISFHVTRGPEITLQPDMPTEQESVSLW
CTADRSTFENLTWYKLGPOPLPIHVGE LPTPVCKNLDLWKLNATMFSNSTNDILIM
5 ELKNASLQDQGDYVCLAQDRKTKKRHCVVVRQLTVLERVAPTITGNLENQTTSIGESI
EVSCITASGNPPPQIMWFKDNETLVEDSGIVLKDGNRNLTI RRV RKEDEGLYCQACSV
LGCAKVEAFFIIEGAQEKTNLEIIILVGTTVIAMFFWLLLVIILGTV. (SEQ.
ID. NO.: 15)

10 8. An expression vector comprising a promoter, and a
DNA sequence encoding a soluble VEGF inhibitor for
expression in recombinant host cells wherein the soluble
VEGF inhibitor is selected from the group consisting of
sVEGF-RI, sVEGF-RII, sVEGF-RTMI and sVEGF-RTMII.
15

9. The expression vector of Claim 8 wherein the DNA
encoding the sVEGF-RI comprises the nucleotide sequence:

20 GCGGACACTCCTCTCGGCTCCTCCCCGCGAGCGCGCGGCTCGGACCGGGCTCCGGGG
CTCGGGTGCAGCGGCCAGCGGGCCTGGCGGCGAGGATTACCGGGGAAGTGGTTGTCTC
CTGGCTGGAGCCGCCAGACGGGCGCTCAGGGCCGGGGCGGGCGGGCGGAACGAGAGG
25 ACGGACTCTGGCGGCCGGGTGCTTGCGCGGGGAGCGCGGCCACCGGGCGAGCAGGCCG
30

- 94 -

CGTCGGGCTCACC ATG GTC AGC TAC TGG GAC ACC GGG GTC CTG CTG
TGC GCG CTG CTC AGC TGT CTG CTT CTC ACA GGA TCT AGT TCA GGT
5 TCA AAA TTA AAA GAT CCT GAA CTG AGT TTA AAA GGC ACC CAG CAC
ATC ATG CAA GCA GGC CAG ACA CTG CAT CTC CAA TGC AGG GGG GAA
10 GCA GCC CAT AAA TGG TCT TTG CCT GAA ATG GTG AGT AAG GAA AGC
GAA AGG CTG AGC ATA ACT AAA TCT GCC TGT GGA AGA AAT GCC AAA
CAA TTC TGC AGT ACT TTA ACC TTG AAC ACA GCT CAA GCA AAC CAC
15 ACT GGC TTC TAC AGC TGC AAA TAT CTA GCT GTA CCT ACT TCA AAG
AAG AAG GAA ACA GAA TCT GCA ATC TAT ATA TTT ATT ACT GAT ACA
20 GGT AGA CCT TTC GTA GAG ATG TAC AGT GAA ATC CCC GAA ATT ATA

25

30

- 95 -

CAC ATG ACT GAA GGA AGG GAG CTC GTC ATT CCC TGC CGG GTT ACG
TCA CCT AAC ATC ACT GTT ACT TTA AAA AAG TTT CCA CTT GAC ACT
5 TTG ATC CCT GAT GGA AAA CGC ATA ATC TGG GAC AGT AGA AAG GGC
TTC ATC ATA TCA AAT GCA ACG TAC AAA GAA ATA GGG CTT CTG ACC
10 TGT GAA GCA ACA GTC AAT GGG CAT TTG TAT AAG ACA AAC TAT CTC
ACA CAT CGA CAA ACC AAT ACA ATC ATA GAT GTC CAA ATA AGC ACA
CCA CGC CCA GTC AAA TTA CTT AGA GGC CAT ACT CTT GTC CTC AAT
15 TGT ACT GCT ACC ACT CCC TTG AAC ACG AGA GTT CAA ATG ACC TGG
AGT TAC CCT GAT GAA AAA AAT AAG AGA GCT TCC GTA AGG CGA CGA
20 ATT GAC CAA AGC AAT TCC CAT GCC AAC ATA TTC TAC AGT GTT CTT

25

30

- 96 -

ACT ATT GAC AAA ATG CAG AAC AAA GAC AAA GGA CTT TAT ACT TGT
CGT GTA AGG AGT GGA CCA TCA TTC AAA TCT GTT AAC ACC TCA GTG
5 CAT ATA TAT GAT AAA GCA TTC ATC ACT GTG AAA CAT CGA AAA CAG
CAG GTG CTT GAA ACC GTA GCT GGC AAG CGG TCT TAC CGG CTC TCT
10 ATG AAA GTG AAG GCA TTT CCC TCG CCG GAA GTT GTA TGG TTA AAA
GAT GGG TTA CCT GCG ACT GAG AAA TCT GCT CGC TAT TTG ACT CGT
GGC TAC TCG TTA ATT ATC AAG GAC GTA ACT GAA GAG GAT GCA GGG
15 AAT TAT ACA ATC TTG CTG AGC ATA AAA CAG TCA AAT GTG TTT AAA
AAC CTC ACT GCC ACT CTA ATT GTC AAT GTG AAA CCC CAG ATT TAC
20 GAA AAG GCC GTG TCA TCG TTT CCA GAC CCG GCT CTC TAC CCA CTG

25

30

- 97 -

GGC AGC AGA CAA ATC CTG ACT TGT ACC GCA TAT GGT ATC CCT CAA
CCT ACA ATC AAG TGG TTC TGG CAC CCC TGT AAC CAT AAT CAT TCC
5 GAA GCA AGG TGT GAC TTT TGT TCC AAT AAT GAA GAG TCC TTT ATC
CTG GAT GCT GAC AGC AAC ATG GGA AAC AGA ATT GAG AGC ATC ACT
10 CAG CGC ATG GCA ATA ATA GAA GGA AAG AAT AAG ATG GCT AGC ACC
TTG GTT GTG GCT GAC TCT AGA ATT TCT GGA ATC TAC ATT TGC ATA
GCT TCC AAT AAA GTT GGG ACT GTG GGA AGA AAC ATA AGC TTT TAT
15 ATC ACA GAT GTG CCA AAT GGG TTT CAT GTT AAC TTG GAA AAA ATG
CCG ACG GAA GGA GAG GAC CTG AAA CTG TCT TGC ACA GTT AAC AAG
20 TTC TTA TAC AGA GAC GTT ACT TGG ATT TTA CTG CCG ACA GTT AAT

25

30

- 98 -

AAC AGA ACA ATG CAC TAC AGT ATT AGC AAG CAA AAA ATG GCC ATC
ACT AAG GAG CAC TCC ATC ACT CTT AAT CTT ACC ATC ATG AAT GTT
5 TCC CTG CAA GAT TCA GGC ACC TAT GCC TGC AGA GCC AGG AAT GTA
TAC ACA CGG GAA GAA ATC CTC CAG AAG AAA GAA ATT ACA ATC AGA
10 GGT GAG CAC TGC AAC AAA AAG GCT GTT TTC TCT CGG ATC TCC AAA
TTT AAA AGC ACA AGG AAT GAT TGT ACC ACA CAA AGT AAT GTA AAA
CAT TAA AGGACTCATTAAAAAGTAACAGTTGTCTCATATCATCTTGATTTATTGTCA
15 CTGTTGCTAACTTTTCAGGCTCGGAGGAGATGCTCCTCCCAAATGAGTTGGGAGATGAT
AGCAGTAATAATGAGACCCCCGGCTCCAGCTCTGGGCCCCCATTCAGGCCGAGGGG

20

25

30

- 99 -

CTGCTCCGGGGGGCCCACTTGGTGACGTTTGGATTGGAGGATCCCTGCACTGCCTTC
TCTGTGTTTGTGCTCTTGCTGTTTTCTCCTGCCTGATAAACAACAACCTGGGATGATC
CTTTCCATTTTATGATGCCAACCTCTTTTTATTTTAAAGCGGCCCCCTATAGT.

5 (SEQ. ID. NO.: 5)

10. The expression vector of Claim 8 wherein the DNA
encoding the sVEGF-RII comprises the nucleotide
sequence:

10

GGTGTGGTCGCTGCGTTTCTCTGCCTGCGCCGGGCATCACTTGCGCGCCGCAGAA
AGTCCGTCTGGCAGCCTGGATATCCTCTCCTACCGGCACCCGCAGACGCCCCCTGCA
GCCGCGGTGCGCGCCCGGGCTCCCTAGCCCTGTGCGCTCAACTGTCTGCGCTGCG
GGGTGCCGCGAGTTCCACCTCCGCGCCTCCTTCTCTAGACAGGCGCTGGGAGAAAG
15 AACCGGTCCCGAGTTCGGGCATTTGCGCCGGCTCGAGGTGCAGGATGCAGAGCAA
GGTGTGCTGCGCCGTGCGCCTGTGGCTCTGCGTGGAGACCCGGGCGCCTCTGTGG
GTTTGCTAGTGTCTCTTGATCTGCCCAGGCTCAGCATACAAAAAGACATACTT
ACAATTAAGGCTAATACAACCTCTTCAAATTACTTGCAGGGGACAGAGGGACTTGGA
CTGGCTTTGGCCCAATAATCAGAGTGGCAGTGAGCAAAGGGTGGAGGTGACTGAGT
20 GCAGCGATGGCCTCTTCTGTAAGACACTCACAATTCCAAAAGTGATCGGAAATGAC
ACTGGAGCCTACAAGTGCTTCTACCGGGAACTGACTTGGCCTCGGTCAATTTATGT
CTATGTTCAAGATTACAGATCTCCATTTATTGCTTCTGTAGTGACCAACATGGAG
TCGTGTACATTACTGAGAACAAAAACAACTGTGGTGATTCCATGTCTCGGGTCC
ATTTCAAATCTCAACGTGTCACTTTGTGCAAGATACCCAGAAAAGAGATTTGTTCC
25 TGATGGTAACAGAATTTCTGGGACAGCAAGAAGGGCTTTACTATTCCAGCTACA
TGATCAGCTATGCTGGCATGGTCTTCTGTGAAGCAAAAATTAATGATGAAAGTTAC
CAGTCTATTATGTACATAGTTGTCGTTGTAGGGTATAGGATTTATGATGTGGTTCT
GAGTCCGTCTCATGGAATTGAACTATCTGTTGGAGAAAAGCTTGTCTTAAATTGTA

30

- 100 -

CAGCAAGAACTGAACTAAATGTGGGGATTGACTTCAACTGGGAATACCCCTTCTTCG
AAGCATCAGCATAAGAACTTGTAACCGAGACCTAAAAACCCAGTCTGGGAGTGA
GATGAAGAAATTTTGTGACACCTTAACTATAGATGGTGTAAACCGGAGTGACCAAG
5 GATTGTACACCTGTGCAGCATCCAGTGGGCTGATGACCAAGAAGAACAGCACATTT
GTCAGGGTCCATGAAAAACCTTTTGTGCTTTTGGAAAGTGGCATGGAATCTCTGGT
GGAAGCCACGGTGGGGGAGCGTGTGAGAATCCCTGCGAAGTACCTTGGTTACCCAC
CCCCAGAAATAAAATGGTATAAAATGGAATACCCCTTGAGTCCAATCACACAATT
AAAGCGGGGCATGTACTGACGATTATGGAAGTGAGTGAAAGAGACACAGGAAATTA
10 CACTGTATCCTTACCAATCCCATTTCAAAGGAGAAGCAGAGCCATGTGGTCTCTC
TGTTGTGTATGTCCACCCAGATTGGTGAGAAATCTCTAATCTCTCTGTGGAT
TCCTACCAGTACGGCACCACTCAAACGCTGACATGTACGGTCTATGCCATTCTCTCC
CCCGCATCACATCCACTGGTATTGGCAGTTGGAGGAAGAGTGCGCCAACGAGCCCA
GCCAAGCTGTCTCAGTGACAAACCCATACCCTTGTGAAGAATGGAGAAGTGTGGAG
15 GACTTCCAGGGAGGAAATAAAATTGCCGTAAATAAAATCAATTTGCTCTAATTGA
AGGAAAAAACAAACTGTAAGTACCCCTTGTATCCAAGCGGCAAATGTGTGAGCTT
TGTAACAATGTGAAGCGGTCAACAAAGTCGGGAGAGGAGAGAGGGTGATCTCCTTC
CACGTGACCAGGGGTCCTGAAATTACTTTGCAACCTGACATGCAGCCCACTGAGCA
GGAGAGCGTGTCTTTGTGGTGCACTGCAGACAGATCTACGTTTGAGAACCTCACAT
20 GGTACAAGCTTGGCCACAGCCTCTGCCAATCCATGTGGGAGAGTTGCCACACCT
GTTTGCAAGAACTTGGATACTCTTTGGAAATTGAATGCCACCATGTTCTCTAATAG
CACAAATGACATTTTGATCATGGAGCTTAAGAATGCATCCTTGCAGGACCAAGGAG
ACTATGTCTGCCTTGCTCAAGACAGGAAGACCAAGAAAAGACATTGCGTGGTCAGG
CAGCTCACAGTCCTAGAGCGTTAA. (SEQ. ID. NO.: 16)

25

11. The expression vector of Claim 8 wherein the DNA encoding the sVEGF-RTM1 comprises the nucleotide sequence:

30 GCGCTCACCATGGTCAGCTACTGGGACACCGGGGTCCTGCTGTGCGGCTGCTCAG
CTGCTGCTTCTCACAGGATCTAGTTCAGGTTCAAAATTTAAAGATCCTGAACTGA

- 101 -

GTTTAAAAGGCACCCAGCACATCATGCAAGCAGGCCAGACACTGCATCTCCAATGC
AGGGGGGAAGCAGCCCATAAATGGTCTTTGCCTGAAATGGTGAGTAAGGAAAGCGA
AAGGCTGAGCATAACTAAATCTGCCTGTGGAAGAAATGGCAAACAATTCTGCAGTA
5 CTTTAACCTTGAACACAGCTCAAGCAAACCACACTGGCTTCTACAGCTGCAAATAT
CTAGCTGTACCTACTTCAAAGAAGAAGGAAACAGAATCTGCAATCTATATATTTAT
TAGTGATACAGGTAGACCTTTTCGTAGAGATGTACAGTGAAATCCCCGAAATTATAC
ACATGACTGAAGGAAGGGAGCTCGTCATTCCCTGCCGGGTACGTCACCTAACATC
ACTGTTACTTTAAAAAAGTTTCCACTTGACACTTTGATCCCTGATGGAAAACGCAT
10 AATCTGGGACAGTAGAAAGGGCTTCATCATATCAAATGCAACGTACAAAGAAATAG
GGCTTCTGACCTGTGAAGCAACAGTCAATGGGCATTTGTATAAGACAACTATCTC
ACACATCGACAAACCAATACAATCATAGATGTCCAAATAAGCACACCACGCCCAGT
CAAATTACTTAGAGGCCATACTCTTGTCCTCAATTGTACTGCTACCACTCCCTTGA
ACACGAGAGTTCAAATGACCTGGAGTTACCCTGATGAAAAAATAAGAGAGCTTCC
15 GTAAGGCGACGAATTGACCAAAGCAATTCCCATGCCAACATATTCTACAGTGTTCT
TACTATTGACAAATGCAGAACAAAGACAAAGGACTTTATACTTGTGCGTGAAGGA
GTGGACCATCATTCAAATCTGTTAACACCTCAGTGCATATATATGATAAAGCATTCT
ATCACTGTGAAACATCGAAAACAGCAGGTGCTTGAAACCGTAGCTGGCAAGCGGTC
TTACCGGCTCTCTATGAAAGTGAAGGCATTTCCCTCGCCGGAAGTTGTATGGTTAA
20 AAGATGGGTTACCTGCGACTGAGAAATCTGCTCGCTATTTGACTCGTGGCTACTCG
TTAATTATCAAGGACGTAAGTGAAGAGGATGCAGGGAATTATACAATCTTGCTGAG
CATAAAACAGTCAAATGTGTTTAAAAACCTCACTGCCACTCTAATTGTCAATGTGA
AACCCAGATTTACGAAAAGGCCGTGTCATCGTTTCCAGACCCGGCTCTCTACCCA
CTGGGCAGCAGACAAATCCTGACTTGTACCGCATATGGTATCCCTCAACCTACAAT
25 CAAGTGGTTCTGGCACCCCTGTAACCATAATCATTCCGAAGCAAGGTGTGACTTTT
GTTCCAATAATGAAGAGTCCTTTATCCTGGATGCTGACAGCAACATGGGAAACAGA
ATTGAGAGCATCACTCAGCGCATGGCAATAATAGAAGGAAAGAATAAGATGGCTAG
CACCTTGGTTGTGGCTGACTCTAGAATTTCTGGAATCTACATTTGCATAGCTTCCA
ATAAAGTTGGGACTGTGGGAAGAAACATAAGCTTTTATATCACAGATGTGCCAAAT
30 GGGTTTCATGTTAACTTGGAATAATGCCGACGGAAGGAGAGGACCTGAAACTGTC
TTGCACAGTTAACAAGTTCTTATACAGAGACGTTACTTGGATTTTACTGCGGACAG

- 102 -

TTAATAACAGAACAATGCACTACAGTATTAGCAAGCAAAAAATGGCCATCACTAAG
GAGCACTCCATCACTCTTAATCTTACCATCATGAATGTTTCCCTGCAAGATTCAGG
CACCTATGCCTGCAGAGCCAGGAATGTATACACAGGGGAAGAAATCCTCCAGAAGA
5 AAGAAATTACAATCAGAGATCAGGAAGCACCATACCTCCTGCGAAACCTCAGTGAT
CACACAGTGGCCATCAGCAGTTCACCACCTTTAGACTGTCATGCTAATGGTGTCCC
CGAGCCTCAGATCACTTGGTTTAAAAACAACCACAAAATACAACAAGAGCCTGGAA
TTATTTTAGGACCAGGAAGCAGCAGCTGTTTATTGAAAGAGTCACAGAAGAGGAT
GAAGGTGTCTATCACTGCAAAGCCACCAACCAGAAGGGCTCTGTGGAAAGTTCAGC
10 ATACCTCACTGTTCAAGGAACCTCGGACAAGTCTAATCTGGAGCTGATCACTCTAA
CATGCACCTGTGTGGCTGCGACTCTCTTCTGGCTCCTATTAACCCTCCTTATCTAA
. (SEQ. ID. NO.: 17)

12. The expression vector of Claim 8 wherein the DNA
15 encoding the sVEGF-RTMII comprises the nucleotide
sequence:

CTCGAGGTGCAGGATGCAGAGCAAGGTGCTGCTGGCCGTCGCCCTGTGGCTCTGCG
TGGAGACCCGGGCCCTCTGTGGGTTTGCCTAGTGTTTCTCTTGATCTGCCCAGG
20 CTCAGCATACAAAAGACATACTTACAATTAAGGCTAATACAACTCTTCAAATTAC
TTGCAGGGGACAGAGGGACTTGGACTGGCTTTGGCCCAATAATCAGAGTGGCAGTG
AGCAAAGGGTGGAGGTGACTGAGTGCAGCGATGGCCTCTTCTGTAAGACACTCACA
ATTCCAAAAGTGATCGGAAATGACACTGGAGCCTACAAGTGCTTCTACCGGGAAAC
TGACTTGGCCTCGGTCAATTTATGTCTATGTTCAAGATTACAGATCTCCATTTATTG
25 CTTCTGTTAGTGACCAACATGGAATCGTGTACATTACTGAGAACAAAAACAAACT
GTGGTGATTCCATGTCTCGGGTCCATTTCAAATCTCAACGTGTCACTTTGTGCAAG
ATACCCAGAAAAGAGATTTGTTCTGATGGTAACAGAATTTCTGGGACAGCAAGA
AGGGCTTTACTATTCCAGCTACATGATCAGCTATGCTGGCATGGTCTTCTGTGAA
GCAAAAATTAATGATGAAAGTTACCAGTCTATTATGTACATAGTTGTGTTGTAGG
30 GTATAGGATTTATGATGTGGTTCTGAGTCCGTCTCATGGAATTGAACTATCTGTTG
GAGAAAAGCTTGTCTTAAATTGTACAGCAAGAACTGAACTAAATGTGGGGATTGAC

- 103 -

TTCAACTGGGAATACCCCTTCTTCGAAGCATCAGCATAAGAACTTGTAACCGAGA
CCTAAAAACCCAGTCTGGGAGTGAGATGAAGAAATTTTTGAGCACCTTAACTATAG
ATGGTGTAACCCGGAGTGACCAAGGATTGTACACCTGTGCAGCATCCAGTGGGCTG
5 ATGACCAAGAAGAACAGCACATTTGTCAGGGTCCATGAAAAACCTTTTGTTGCTTT
TGGAAGTGGCATGGAATCTCTGGTGGAAGCCACGGTGGGGGAGCGTGTGAGAATCC
CTGCGAAGTACCTTGGTTACCCACCCCCAGAAATAAAATGGTATAAAAATGGAATA
CCCCTTGAGTCCAATCACACAATTAAGCGGGGCATGTACTGACGATTATGGAAGT
GAGTGAAAGAGACACAGGAAATTACACTGTCTCCTTACCAATCCCATTTCAAAGG
10 AGAAGCAGAGCCATGTGGTCTCTCTGGTTGTGTATGTCCACCCCAGATTGGTGAG
AAATCTCTAATCTCTCTCTGTGGATTCTTACCAGTACGGCACCCTCAAACGCTGAC
ATGTACGGTCTATGCCATTCTCTCCCCGCATCACATCCACTGGTATTGGCAGTTGG
AGGAAGAGTGCGCCAACGAGCCCAGCCAAGCTGTCTCAGTGACAAACCCATACCCCT
TGTGAAGAATGGAGAAGTGTGGAGGACTTCCAGGGAGGAAATAAAATTGCCGTTAA
15 TAAAAATCAATTTGCTCTAATTGAAGGAAAAAACAAACTGTAAAGTACCCTTGTTA
TCCAAGCGGCAAATGTGTGTCAGCTTTGTACAAATGTGAAGCGGTCAACAAAGTCGGG
AGAGGAGAGAGGGTGATCTCTTCCACGTGACCAGGGGTCTGAAATTACTTTGCA
ACCTGACATGCAGCCCACTGAGCAGGAGAGCGTGTCTTTGTGGTGCACTGCAGACA
GATCTACGTTTGAGAACCTCACATGGTACAAGCTTGGCCACAGCCTCTGCCAATC
20 CATGTGGGAGAGTTGCCCACACCTGTTTGCAAGAACTTGGATACTCTTTGGAAATT
GAATGCCACCATGTTCTCTAATAGCACAAATGACATTTTGATCATGGAGCTTAAGA
ATGCATCCTTGCCAGGACCAAGGAGACTATGTCTGCCTTGCTCAAGACAGGAAGACC
AAGAAAAGACATTGCGTGGTCAGGCAGCTCACAGTCCTAGAGCGTGTGGCACCAC
GATCACAGGAAACCTGGAGAATCAGACGACAAGTATTGGGGAAAGCATCGAAGTCT
25 CATGCACGGCATCTGGGAATCCCCCTCCACAGATCATGTGGTTTAAAGATAATGAG
ACCCCTGTAGAAGACTCAGGCATTGTATTGAAGGATGGGAACCGGAACCTCACTAT
CCGCAGAGTGAGGAAGGAGGACGAAGGCCTCTACACCTGCCAGGCATGCAGTGTC
TTGGCTGTGCAAAAGTGGAGGCATTTTTTCATAATAGAAGGTGCCAGGAAAAGACG
AACTTGGAATCATTATTCTAGTAGGCACGACGGTGATTGCCATGTTCTTCTGGCT
30 ACTTCTTGTATCATCCTAGGGACCGTTTAA. (SEQ. ID. NO.: 18)

- 104 -

13. A recombinant host cell containing the expression vector of Claim 8.

5 14. A method for inhibiting VEGF receptor function comprising the administration of the VEGF inhibitor of Claim 1 in an amount sufficient to inhibit VEGF receptor function.

10 15. The method of Claim 14 wherein the VEGF inhibitor is selected from the group consisting of sVEGF-RI, sVEGF-RII, sVEGF-RTMI, and sVEGF-RTMII.

15 16. A pharmaceutical composition comprising the inhibitor of Claim 1 and a pharmaceutically acceptable carrier.

20 17. The pharmaceutical composition of Claim 16 wherein the inhibitor is selected from the group consisting of sVEGF-RI, sVEGF-RII, sVEGF-RTMI, and sVEGF-RTMII.

25 18. A method for inhibiting angiogenesis comprising the administration of the VEGF inhibitor of Claim 1 in an amount sufficient to inhibit angiogenesis.

30

1/20




FIG. 1

2/20

CGGGACACTCCTCTCGGCTCCTCCCGGCAGCGGGCGGCTCGGAGCGGGCTCCGGGG
CTCGGGTGAGCGGCCAGCGGGCTGGCGGCGAGGATTACCGGGGAAGTGGTTGTCTC
CTGGCTGGAGCCGCGAGACGGCGCTCAGGGCGGGCGGCGGGCGGCGGAACGAGAG
GACGGACTCTGGCGGCCGGTCTGGCCGGGGAGCGGGGACCGGGCGAGCAGGC
CGGTGCGGCTCACCATGGTCAGCTACTGGGACACCGGGGCTGCTGTGCGCGCTGCTC
AGCTGTCTGCTTCTCACAGGATCTAGTTCAGGTTCAAAATTAAGATCCTGAAGTTTA
AAGGCACCCAGCACATGCAAGCAGGCCAGCACACTGCATCTCCAATGCGAGGGGAAG
CAGCCCATAAATGGTCTTTGCCCTGAAATGGTGAGTAAGGAAAGCGAAGGCTGAGCATAACT
AAATCTGCCCTGTGGAAGAAATGGCAACAATTCGCGAGTACTTTAACCTTGAACACAGCTCAA
GCAACCCACACTGGCTTCTACAGCTGCAAAATATCTAGCTGTACCTACTTCAAGAAGAAGGA
AACAGAACTCTGCAATCTATATATTTATTAGTGATACAGGTAGACCTTTCTGATAGATGTACAG
TGAAATCCCCGAAATTATACACATGACTGAAGGAAGGAGCTCGTCAATCCCTGCCGGGTTA
CGTCACCTAACATCACTGTACTTTAAAGTTTCCACTTGACACTTTGATCCCTGATGGAA
AACGCATAATCTGGGACAGTAGAAGGCTCAATGGGCATTGTATAGACAACTATCTCACACA
GGGCTTCTGACCTGTGAAGCAACAGTCAATGGGCATTGTATAGACAACTATCTCACACA
TCGACAAACCAATACAATCATAGATGTCCAAATAGCACACACCCAGTCAAAATTAAGT
AGGCCATACTCTTGCTCTCAATTTGCTGCTACCACTCCCTTGAACACGAGTTCAAAATGAC
CTGGAGTTACCTGATGAATAAATAAGAGAGCTTCCGTAAAGGCGACGAAATGACCAAGCA
ATTCCCATGCCAACAATTTCTACAGTGTCTTACTATTGACAAATGCAGAAACAAAGACAAG
GACTTTTACTTGTGCGTGAAGGAGTGGACCATCAATCAAAATCTGTTAACACCTCAGTGCATA
TATATGATAAAGCATTCACTGTAAGACATCGAAACAGCAGGTGCTTGAACCGTAGCT
GGCAAGCGGTCTTACCGGCTCTCTATGAAGTGAAGGCATTTCCCTCGCCGGAAGTTGTAT

FIG. 2A

SUBSTITUTE SHEET (RULE 26)

3/20

GGTTAAAGATGGGTTACCTGCGACTGAGAAATCTGCTCGCTATTTGACTCGTGGCTACTCG
TTAATTATCAAGBACGTAACTGAAGAGGATGCGAGGAATTATACAATCTTGCTGAGCATAAA
CAGTCAATGTGTTAAACCGTCACTGCCACTCTAATGTCAATGTGAACCCAGATTAC
GAAAGGCGGTGTCATCGTTCCAGACCGGCTCTACCCACTGGGAGCAGACAAATCC
TGACTTGTAACCGCATATGGTATCCCTCAACCTACAATCAAGTGGTTCTGGCACCCCTGTAAAC
CATAATCATTCGGAAGCAAGGTGTGACTTTTGTCCAAATATGAAGAGTCCTTTATCCTGGAT
GCTGACAGCAACATGGGAACAGAAATTGAGAGCATCACTCAGCGCATGGCAATAATAGAAG
GAAAGAAATAGATGGCTAGCACCTTGGTTGTGGCTGACTCTAGAAATTTCTGGAATCTACATTT
GCATAGCTTCCAATAAGTTGGGACTGTGGGAAGAAACATAAGCTTTTATATCACAGATGTG
CCAAATGGGTTTCATGTTAACTTGGAAAAATGCCGACGGAGGAGGACCTGAACTGTC
TTGCACAGTTAACAGTTCTTATACAGAGACGTTACTTGGATTTTACTGCGGACAGTTAATAA
CAGAACAAATGCACACAGTATTAGCAAGCAAAAATGGCCATCACTAAGGAGCACTCCATCA
CTCTTAATCTTACCATCATGAATGTTTCCCTGCAAGATTCAGGCACCTATGCCCTGCAGAGCCA
GGAATGTATACACAGGGGAAGAAATCCTCCAGAAAGAAATTAACAATCAGAGGTGAGCAC
TGCAACAAAAAGGCTGTTTCTCTCGGATCTCCAAATTTAAAGCAAGGAATGATTGTACC
ACACAAGTAATGTAAACATTAAGGACTCATTAAAGTAACAGTTGTCTCATATCATCTTG
ATTTATTGTCACCTGTTGCTAACTTTCAGGCTCGGAGGAGATGCTCCTCCAAATAGTTTG
GAGATGATAGCAGTAATAATGAGACCCCGGCTCCAGCTCTGGGCCCCCATTGAGGCCG
AGGGGGCTGCTCCGGGGGCCGACTTGGTGACGTTTGGATTGGAGGATCCCTGCACTG
CCTTCTCTGTTTGTGCTCTTGTCTGTTTCTCTGCTGATAAACAACAACTTGGGATGAT
CCTTTCCATTTTGATGCCAACCTCTTTTATTTTAAAGCGGCGCCTATAGT
(SEQ. ID. NO.: 5)

FIG. 2B

SUBSTITUTE SHEET (RULE 26)

4/20

MVSYWDTGVLLCALLSCLLLTGSSSGSKLDPELSUKGTQHIMQAGQTLHLQC
RGEAAHKWSLPEMVSKESERLSITKSACGRNGKQFCSTLTNTAQANHTGFYS
CKYLAVPTSKKKETESAIYIFISDTGRPFVEMYSEIPEIHMTEGRELVIPCRVTSP
NITVTLKFPDLTLPDGGKRIIWDNRKGFISNATYKEIGLLTCEATVNGHLYKTNLY
THRQNTIIDVQISTPRPVKLLRGHTLVNCTATTPLNTRVQMTWSYPDEKNKR
ASVRRRIDQSNSHANIFYSVLTIDKMQNDKGLYTCRVRSGPSFKSVNTSVHIY
DKAFITVKHRKQQVLETVAGKRSYRLSMKVKAFFSPVWVWLKDGLPATEKSAR
YLTRGYSIIKDVTEEDAGNYTILLSIKQSNVFNLTATLVNVKPKIYEKAVSSFP
DPALYPLGSRQILTCTAYGIPQPTIKWFWHPCHNHNHSEARCDFCSSNNEESFILD
ADSNMGNRIESITQRMALIEGKNKMASTLVWADSRISGIYICIASNKVGTVGRNISF
YITDVPNGFHVNLKEMPTGEDLKLSCVTNKFYRDVTWILLRTVNNRTMHYSIS
KQKMAITKEHSITLNLTIMNVSLQDSGTYACRARNVYTGEIILQKKEITIRGEHCN
KKAVFSRISKFKSTRNDCTTQSNVKH (SEQ. ID. NO.: 6)

FIG. 3

5/20




FIG. 4

6/20





FIG. 5

7/20

**FIG. 6**




FIG. 7A




FIG. 7B




FIG. 8




FIG. 9

SUBSTITUTE SHEET (RULE 26)

10/20

GGTGTGGTCGCTGCGTTTCCTCTGCCCTGCGCCGGGCATCACTTGCGGCCCGCAGAAAGTC
CGTCTGGCAGCCTGGATATCCTCTCTACCGGCACCCGACGCGCCCTGCAGCCGCGGT
CGGCGCCCGGCTCCCTAGCCCTGTGCGTCAACTGTCTGCGCTGCGGGTGCCGCGAG
TTCACCTCCGCGCTCTCTCTAGACAGGCGCTGGAGAAAGAACCGGCTCCCGAGTTC
CGGCATTTCCGCCCGCTCGAGGTGCAGGATGCAGAGCAAGGTGCTGCGCGTCCGCGCT
GTGGCTCTGCGTGGAGACCCGGCGCCTCTGTGGGTTTGCCTAGTGTCTCTTGATCTG
CCCAGGCTCAGCATACAAAGACATACATAATTAAAGCTAATACACTCTTCAAATTACT
TGCAGGGGACAGAGGACTTGACTGGCTTTGGCCCAATAATCAGAGTGGCAGTGAGCAAA
GGGTGGAGGTGACTGAGTGCAGCGATGGCTCTTCTGTAAAGACACTCACAATTCCAAAGT
GATCGGAAATGACACTGGAGCCTACAAGTGCTTCTACCGGAAACTGACTTGGCCTCGGTC
ATTTATGTCTATGTTCAAGATTACAGATCTCCATTTATTGCTTCTGTAGTGACCAACATGGAG
TCGTGTACATTACTGAGAACAAACAAACTGTGGTGATTCATGTCTCGGTCATTTCAA
ATCTCAACGTGTCACTTTGTGCAAGATACCCAGAAAGAGATTGTTCCGTGATGTAACAGAA
TTTCTGGGACAGCAAGAAAGGCTTTACTATTCCAGCTACATGATCAGCTATGCTGGCATG
GTCTTCTGTGAAGCAAAATTAATGATGAAAGTTACCAAGTCTATTATGTACATAGTTGTCGT
GTAGGGTATAGGATTTATGATGTGGTTCTGAGTCCGTCATGGAATTGAACATCTGTGGA
GAAAAGCTTGCTTAAATTGTACAGCAAGAACTGAACATAATGTGGGATTGACTTCAACTGG
GAATACCCCTTCTCGAAGCATCAGCATAAGAACTTGTAAACCGAGACCTAAACCCAGTCT
GGGAGTGAGATGAAGAAATTTTGTAGCACCTTAACATAGATGGTGTAAACCGAGTGACCA

FIG. 10A

11/20

AGGATTGTACACCTGTGCAGCATCCAGTGGGTGATGACCAAGAAGAACAGCACATTGTGCA
GGGTCCATGAAAAACCTTTTGTGCTTTTGGAGTGGCATGGAATCTCTGGTGGAAAGCCACG
GTGGGGAGCGTGTACAGATCCCTGCGAAGTACCTTGGTTACCCACCCCGAGAAATAAAT
GGTATAAAAAATGGAATACCCCTTGAGTCCAAATCACACAAATTAAGCGGGGCATGTACTGACG
ATTATGGAAGTGAGTGAAGAGACACAGGAAATTACACTGTCTCCTTACCAATCCCATTCA
AAGGAGAAGCAGAGCCATGTGCTCTCTGTTGTGTATGTCCACCCAGATTGGTGAGA
AATCTCTAATCTCTCTGTTGATTCTTACAGTACGGCACCCACTCAACGCTGACATGTACG
GTCTATGCCATTCTCCCGCATCACATCCACTGGTATGGCAGTTGGAGGAAGAGTGCG
CCAACGAGCCCGCAAGCTGTCTCAGTGACAAACCCATACCCCTTGTGAAGATGGAGAAG
TGTGGAGGACTTCCAGGGAGGAAATAAAATTGCCGTTAATAAAAAATCAATTTGCTCTAAATTGA
AGGAAAAAACAAACTGTAACTACCTTGTATCCAAGCGGCAATGTGTCAGCTTTGTACAA
ATGTGAAGCGGTCAACAAAGTCGGGAGAGGAGAGGGTGATCTCCTTCCACGTGACCAGG
GGTCCTGAAATTACTTTGCAACCTGACATGCAGCCCATCATGTTGACAGCGTGTCTTTGTG
GTGCACTGCAGACAGATCTACGTTTGAGAACCTCACATGTTGCAAGCTTGGCCACAGCCTC
TGCCAAATCCATGTGGGAGATTGCCACACCTGTTTGCAGAACTTGGATACTCTTTGGAA
TTGAATGCCACCATGTTCTCTAATAGCACAAATGACATTTGATCATGGAGCTTAAGAAATGCA
TCCTTGCAAGGACCAAGGAGACTATGTCTGCTTCAAGACAGGAAAGACCAAGAAAGAC
ATTGCGTGGTCAAGGCAGCTCACAGTCTAGAGCGTTAA (SEQ. ID. NO.: 16)

FIG. 10B

12/20

MQSKVLLAVALWLCVETRAASVGLPSVSLDLPRLSIQKDILTIKANTTLQITCRGQ
RDLDWLWPNNQSGSEQRVEVTECSDGLFCKLTLPKVIGNDTGAYKCFYRETD
LASVIYVQDYRSPFIASVSDQHGVVYITENKNKTVPCLGSISNLNVSLCARY
PEKRFVPDGNRISWDSKKGFTIPSYMISYAGMVCFCEAKINDESYQSIMYIVVVVG
YRIYDVLSPSHGIELSVGEKLVNCTARTELVNIGIDFNWEYPSSKHQHKLLVN
RDLKTQSGSEMKKFLSTLTIDGVTRSDQGLYTCAASSGLMTKKNSTFVRVHEK
PFVAFGSGMESLVEATVGERVRIPAKYLGYPPEIKWYKNGIPLESNHTIKAGHV
LTIMEVSRDGTNYTVILTNPISKEKQSHVSVLVVYVPPQIGEKSLISPVDYQYG
TTQTLTCTVYAIPPPHHIHWYQLEEECANEPSQAVSVTNYPCEEWRSEDF
QGGNKIYVKNQFALIEGKNKTSTLVIAQANVSALYKCEAVNKVGRGERVISFH
VTRGPEITLQPDQMPTQESVSLWCTADRSTFENLTWYKLGPOPLPIHVGEPLPT
PVCKNLDTLWKLNATMFSNSTNDILIMELKNASLQDQGDYVCLAQDRKTKKRH
CWRQLTVLER... (SEQ. ID. NO.: 13)

7
FIG. 11

13/20

GGTGTGGTCGCTGCGTTTCTCTGCTGCGCGGGCATCACTTGCGCGCCGAGAAAGTC
CGTCTGGCAGCCTGGATATCCTCTCTACCGGCACCCGAGACGCCCTGCAGCCGCGGT
CGGCGCCCGGGTCCCTAGCCTGCTCACTGCTGCGCTGCGGTGCGGGTGCGCGGAG
TTCCACCTCCGCGCTCCTTCTCTAGACAGCGCTGGAGAAAGAACCGCTCCGAGTTC
CGGCATTTGCGCCGGCTCGAGGTGCAGGTGACAGAGCAAGTGCTGCTGGCGTCCGCCCT
GTGGCTCTGCGTGGAGACCCGGCGCCCTCTGTGGGTTTGCTAGTGTCTCTGATCTG
CCCAGGCTCAGCATACAAAGACATAGTTACAATTAGGCTAATACAACCTCTTCAATTACT
TGCAGGGGACAGAGGACTTGGACTGGCTTTGGCCCAATAATCAGAGTGGCAGTGAGCAAA
GGGTGGAGGTGACTGAGTGCAGCGATGGCTCTTCTGTAGACACTCACAATCCAAAAGT
GATCGGAAATGACACTGGAGCCTACAAGTCTTCTACCGGAAACTGACTTGGCCTCGGTC
ATTTATGTCTATGTTCAAGATTACAGATCTCCATTTATTGCTTCTGTTAGTGACCAACATGGAG
TCGTGTACATTACTGAGAACAAACAAACTGTGGTGATTCATGTCTCGGGTCCATTTCAA
ATCTCAACGTGTCACTTTGTGCAAGATACCCAGAAAGAGATTGTTCTGTGTAACAGAA
TTTCTGGGACAGCAAGAGGCTTTACTATTCAGCTACATGATCAGCTATGCTGGCATG
GTCTTCTGTGAAGCAAAATTAAATGATGAAGTTACCAGTCTATTATGTACATAGTTGTCGTT
GTAGGGTATAGGATTTATGATGTGGTTCTGAGTCCGTCTCATGGAATTGAACATCTGTGGA
GAAAAGCTTGTCTTAAATTGTACAGCAAGAACTGAACATAATGTGGGATTGACTTCAACTGG
GAATACCTTCTTCGAAGCATCAGCATAAGAACTGTAAACCGAGACCTAAAACCCAGTCT
GGGAGTGAGATGAAGAAATTTTGGAGCACCTTAACATATAGATGGTGTAAACCGGAGTGACCA
AGGATTGTACACCTGTGCAGCATCCAGTGGGCTGATGACCAAGAAACAGACATTTGTCA
GGGTCCATGAAAACCTTTTGTGCTTTTGGAAAGTGGCATGGAATCTCTGGTGGAAAGCCACG
GTGGGGAGCGTGTCAAGATCCCTGCGAAGTACCTTGGTTACCCACCCCGAGAAATAAAAT

FIG. 12A

14/20

GGTATAAAATGGAATACCCCTTGAGTCCAATCACAAATTAAGCGGGGCATGTACTGACG
ATTATGGAAGTGAGTGAAGAGACACAGGAATACACTGTCACTCTTACCAATCCCATTTCA
AAGGAGAAGCAGAGCCATGTGGTCTCTGGTTGTATGTCCCACCCAGATTGGTGAGA
AATCTCTAATCTCTCTGTGATTCTACCACTACGCCACCACTCAAACGCTGACATGTACG
GTCTATGCCATTCTCCCCGATCACATCCAGTGTATTGGCAGTTGGAGGAAGATGGCG
CCAAAGAGCCAGCCAGCTGTCTCAGTGACAAACCATACCCCTTGTGAAGATGGAGAAG
TGTTGGAGGACTCCAGGGAGGAATAAATGCCGTTAATAAAATCAATTTGCTCTAAATTGA
AGGAAAACAAACACTGTAGTACCTTGTTATCCAAGCGCAATGTGTACGCTTTGTACAA
ATGTGAAGCGGTCAACAAGTCGGAGAGGAGAGGGTGATCTCCTCCACGTGACCAAG
GGTCTCTGAAATTACTTTGCAACCTGACATGCAGCCACTGAGCAGGAGCGTGCTTTTGTG
GTSCACTGCAGACAGATCTAGTTTGAGAACCTGCATGGTACAAGCTTGGCCACAGCCCT
TGCCAAATCCATGTGGGAGATTGCCACACCTGTTGCAAGAACTTGGATACTCTTTGGAA
TTGAATGCCACCATGTTCTCTAATAGCACAAATGACATTTGATCATGGAGCTTAAGAATGCA
TCCTTGCAGGACCAAGGAGACTATGTCTGCTTGTCTCAAGCAGGAAGACCAAGAAAGAC
ATTGCGTGGTCAGGCACTCACAGTCTAGAGCGTGTGGCACCCACGATCACAGGAACCT
GGAGAAATCAGACGACAAGTATTGGGAAAGCATCGAAGTCTCATGCAAGCATCTGGGAAT
CCCCCTCCACAGATCATGTGGTTTAAAGATAATGAGACCCCTTGTAGAAAGACTCAGGCATTGT
ATTGAAGGATGGGAACCGGAACCTCACTATCCGCAGAGTGAGGAAGGAGGACGAAGGCCT
CTACACCTGCCAGGCATGCAGTGTCTTGGCTGTGCAAAAGTGGAGGCATTTTTCATAATAG
AAAGTGGCCAGGAAAGCAGAACTTGGAAATCATATTCTAGTAGGCACGACGGTATTGCC
ATGTTCTCTGGCTACTTCTTGTCATCATCCTAGGGACCGTTTAA (SEQ. ID. NO.: 18)

FIG. 12B

15/20

MQSKVLLAVALWLCVETRAASVGLPSVSLDLPRLSIQKDILTIKANTTLOITCRGQ
RDLDWLWPNNQSGSEQRVEVTECSDGLFCKLTIPKVIGNDTGAYKCFYRETD
LASVIYVYVQDYRSPFIASVSDQHGVVYITENKNKTVVIPCLGSISNLNVSLCARY
PEKRFVPDGNRISWDSKKGFTIPSYMISYAGMVCFEAKINDESYQSIMYIVVVVG
YRIYDVVLSPSHGIELSVGEKLVNCTARTELVGIDFNWEYPSSKHQHKLVN
RDLKTQSGSEMKKFLSTLTIDGVTRSDQGLYTCAASSGLMTKKNSTFVRVHEK
PFVAFSGMESLVEATVGERVRIPAKYLGYPPEIKWYKNGIPLESNHTIKAGHV
LTIMEVSEKDTGNYTVILTNPISKEKQSHVVSLVYVPPQIGEKSLISPVDSYQYG
TTQTLTCTVYAIPPPHHIHWYWQLEEECANEPSQAVSVTNPYPCEEWRSEDF
QGGNKIAVNKNQFALIEGKNKTVSTLVIQAANVSALYKCEAVNKVGRGERVISFH
VTRGPEITLQPDMPTEQESVSLWCTADRSTFENLTWYKLGPOPLPIHVGEPT
PVCKNLDTLWKLNATMFSNSTNDILIMELKNASLQDQGDYVCLAQDRKTKKRH
CVVRQLTVLERVAPTITGNLENQTTSIGESIEVSCTASGNPPPQIMWFKDNETLV
EDSGIVLKDGNRNLTIRRVKEDGLYTQACSVLGCAKVEAFFIIEGAQKTNL
EIIILVGTTVIAMFFWLLLVIILGTV... (SEQ. ID. NO.: 15)

FIG. 13

SUBSTITUTE SHEET (RULE 26)

16/20

GGGCTACCATGGTCAGCTACTGGACACCGGGGTCCTGCTGTGCGCGTCTCAGCTGT
CTGCTTCTCAGAGATCTAGTTCAGGTTCAAAATTAAAGATCCTGAACCTGAGTTAAAGGC
ACCCAGCACATCATGCAAGCAGGCCAGACACTGCATCTCCAATGCAGGGGGAAGCAGCC
CATAAATGGTCTTTGCCGTAATGGTGAGTAAGGAAAGCGAAAGGCTGAGCATAACTAAATC
TGCCCTGTGGAAGAAATGGCAACAATTCTGCAGTACTTTAACCTTGAACACAGCTCAAGCAA
ACCACACTGGCTTCTACAGCTGCAATATCTAGCTGACAGTACCTTTCAAGAAGAAGGAAACA
GAATCTGCAATCTATATATTTATTAGTGATACAGGTAGACCTTTGTAGAGATGTACAGTGAA
ATCCCCGAAATTATACACATGACTGAAGGAAGGAGCTCGTCAATCCCTGCCGGTTACGTC
ACCTAACATCACTGTTACTTTAAAAAGTTTCCACTTGACACTTTGATCCCTGATGBAAACG
CATAATCTGGACAGTAGAAAGGGCTTCATCATATCAATGCAATGCAACGTACAAAGAAATAGGGC
TTCTGACCTGTGAAGCAACAGTCAATGGCATTTGTATAAGACAAACTATCTCACACATCGAC
AAACCAATACAATCATAGATGTCCAAATAAGCACACACGCCAGTCAAAATTACTTAGAGGC
CATACTCTTGTCCCTCAATTGTACTGCTACCACTCCCTTGAACACGAGAGTTCAAAATGACCTGG
AGTTACCCCTGATGAAAAAATAAGAGAGCTTCCGTAAGGCGACGAA TTGACCAAAGCAATTC
CCATGCCAACATATTTCTACAGTGTCTTACTATTGACAAATGCAAAACAAAGACAAAGGACT
TTATACTTGTGCGTGAAGGAGTGGACCATCATTTCAAACTGTTAACACCTCAGTGCATATATA
TGATAAAGCAATTCATCACTGTGAACATCGAAACAGCAGGTGCTTGAACCCGTAGCTGGCA
AGCGGTCTTACCGGCTCTCTATGAAAGTGAAGGCATTTCCCTCGCCGGAAGTTGTATGGTTA
AAAGATGGGTTACCTGCGACTGAGAAATCTGCTCGCTATTGACTCGTGGCTACTCGTTAAT

FIG. 14A

17/20

TATCAAGGACGTAACTGAAGAGGATGCAGGGAATTATACAATCTTGGTGAGCATAAACAGT
CAAATGTGTTTAAACCTCACTGCCACTCTAATTGTCAATGTGAACCCAGATTTACGAAA
AGCGGTGTCATCGTTCCAGACCCGGCTCTCTACCCACTGGGAGCAGACAAATCCTGAC
TTGTACCGCATATGGTATCCCTCAACCTACAATCAAGTGGTTCTGGCACCCCTGTAAACATAA
TCATTCGGAAGCAAGGTGTGACTTTTGTTCCTAATAATGAAGAGTCCCTTTATCCTGGATGCTGA
CAGCAACATGGGAACAGAAATTGAGAGCATCACTCAGCGCATGGCAATAATAGAAGGAAAG
AATAAGATGGCTAGCACCTTGGTTGGCTGACTCTAGAAATTTCTGGAATCTACATTTGCATA
GCTTCCAATAAAGTTGGGACTGTGGGAAGAAACATAAGCTTTTATATCACAGATGTGCCAAAT
GGGTTTCATGTAACTTGGAAATAATGCCGACGGAAGGAGGACCTGAAACTGTCTTGAC
AGTTAACAAAGTTCTTATACAGAGACGTTACTTGGATTTTACTCGGACAGTTAAATACAGAAC
AATGCACCTACAGTATTAGCAAGCAAAATAATGGCCATCACTAAGGAGCACTCCATCCTTTAA
TCTTACCATCATGAATGTTTCCCTGCAAGATTCAGGCACCTATGCCTGCAGAGCCAGGAATG
TATACACAGGGGAAGAAATCCTCCAGAAGAAAGAAATTACAATCAGAGATCAGGAAGCACCA
TACCTCCTGCGAAACCTCAGTGATCACACAGTGGCCATCAGCAGTCCACCACCTTAGACTG
TCATGCTAATGGTGTCCTCCGAGCCTCAGATCAGTCTGGTTTAAACCAACCAAAAATACAACA
AGAGCCTGGAATATTATAGGACCAGGAAGCAGCAGCTGTTTATTGAAAGAGTCACAGAAG
AGGATGAAGGTGTCATCACTGCAAGCCACCACCAAGAGGCTCTGTGGAAAGTTTCAGC
ATACCTCACTGTTCAAGGAACCTCGGACAAGTCTAATCTGGAGCTGATCACTTAACATGCA
CCTGTGGCTGCGACTCTCTTCTGGCTCCTATTAAACCCTCCTTATCTAA (SEQ. ID. NO.: 17)

FIG. 14B

SUBSTITUTE SHEET (RULE 26)

18/20

MVSYWDTGVLLCALLSCLLLTGSSSGSKLDPELSLKGTHIMQAGQTLHLQC
RGEAAHKWSLPPEMVSKESERLSITKSACGRNGKQFCSTLTLNTAQANHTGFYS
CKYLAVPTSKKKESESAYIFISDTGRPFVEMYSEIPEIHMTEGRELVIPCRVTSP
NITVTLKKFPLDTLIPDGKRIIWDSRKGFIISNATYKEIGLLTCEATVNGHLYKTNYL
THRQNTNIIDVQISTPRPVKLLRGHTLVLNCTATTPLNTRVQMTWSYPDEKNKR
ASVRRRIDQSNHANIFYSVLTIDKMQNKDKGLYTCRVRSGPSFKSVNTSVHIY
DKAFITVKHRKQQVLETVAGKRSYRLSMKVKAFPSPEVWVWLDGLPATEKSAR
YLTRGYSLLIKDVTEEDAGNYTILLSIKQSNVFNLTATLIVNVKPOIYEKAVSSFP
DPALYPLGSRQILTCTAYGIPQPTIKWFWHPCNHNHSEARCDFCNNEESFILD
ADSNMGNRIESITORMAIIIEGKNKMASTLVVADSRISGIYICIASNKVGTVGRNISF
YITDVPNGFHVNLKMPTEGEDLKLSTVKNFLYRDVTWILLRTVNNRTMHYSIS
KQKMAITKEHSITLNLTIMNVSLQDSGTACRARNVYTGEELQKKEITIRDQEAP
YLLRNLSDHTVAISSSTLDCHANGVPEPQITWFKNNHKIQQEPGIIILGPGSSSTLF
IERVTEEDEGVYHCKATNQKGSVESSAYLTVQGTSDKSNLELITLTCTCVAATLF
WLLLTLLI (SEQ. ID. NO.:14)

FIG. 15

SUBSTITUTE SHEET (RULE 26)

19/20




FIG. 16

20/20




FIG. 17

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/01957

A. CLASSIFICATION OF SUBJECT MATTER

IPC(S) : C07K 13/00; C12P 21/00; C12N 5/00, 15/00

US CL : 435/69.1, 240.1, 320.1; 530/350; 536/23.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 240.1, 320.1; 530/350; 536/23.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, Medline, Biosis, WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Journal of Cellular Physiology, Volume 149, Number 1, issued October 1991, Bikfalvi et al, "Interaction of Vasculotropin/Vascular Endothelial Cell Growth Factor with Human Umbilical Vein Endothelial Cells: Binding, Internalization, Degradation, and Biological Effects", pages 50-59, see abstract.	1
Y		14, 15, 18
Y	Science, Volume 255, issued 21 February 1992, De Vries et al, "The fms-Like Tyrosine Kinase, a Receptor for Vascular endothelial Growth Factor", pages 989-991, see abstract and fig. 1.	1-18

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

12 MAY 1994

Date of mailing of the international search report

JUN 03 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Sally P. Teng

Telephone No. (703) 308-0196

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Oncogene, Volume 5, issued 1990, Shibuya et al, "Nucleotide Sequence and Expression of a Novel Human Receptor-Type Tyrosine Kinase Gene (flt) Closely Related to the fms Family", pages 519-524, see abstract and page 521.	1-18
Y	Biochemical and Biophysical Research Communications, Volume 187, Number 3, issued 30 September 1992, Terman et al, "Identification of the KDR Tyrosine Kinase as a Receptor for Vascular Endothelial Cell Growth Factor", pages 1579-1586, see summary and page 1583.	1-18